
Oils" in	nto Inert Ingredient 14sgint.	PM Pre	e issued: 1/18/90 (Green) (v. Date Issued: 7/14/89	
(9-Squa	re Logo ⁽⁾)	FROMT PANEL		
		PURINA	EPA Reg. No. 602-187	
		BRAND		
		DAIRY SPRAY		
		Ho Flies, Mosquitoes, an		
	INGREDIENTS:			
	ethrins eronyl butoxide, technical *			
INERT IN	IGREDIENTS	••••••••••••••••••••••	<u>99.46</u> % 100.00%	
	sts of 0.384% of (Butylcarbit 6 of Related Compounds.	yl) (6-Propylpiperonyl) Ether and	
Inerts i	include Aliphatic petroleum h	ydrocarbons		
	KEF.P	OUT OF REACH OF CHILDR	EN	
		CAUTION		
	SEE OTH	ER WARNINGS ON BACK PA	NEL	
NET 5 GAL		D-2497		
(Note:	Type size and prominence will conform with requirements for precautionary labeling.)			
		Manufactured By	•••	

PURINA MILLS, INC. P.O. Box 66812 St. Louis, Missouri 63166-6812 U.S.A.



Not Reviewed - Registrate the ns to be in accordance with EVE grade to - Prestored - Product Guidance for Phase 2 Prestored - Page 2-12, Active to Inert Change in Status.

3

么

Back Panel

How To Use Purina Dairy Spray BRAND

PURINA Dairy Spray is a quick knockdown contact spray for killing and repelling horn flies, house flies, mosquitoes and gnats and also kills lice. For fly control, apply daily; after using 3 days the residual repellent and killing action may be sufficient so that applications may be needed only every other day.

Dairy Cattle:

<u>Fly Control</u> - Apply Purina Dairy Spray as a mist to the hair coat. Hold sprayer about 18 inches from the cow; apply mist to entire body from mid-neck region to tail, including the underline and the legs. The amount of spray used should not exceed 1 to 2 ounces per adult animal. Spray calves more sparingly. Do not wet the skin. Wash udder before milking.

<u>Lice Control</u> - Spray the animals with 2 ounces of Purina Dairy Spray. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks.

<u>Dairy Barn and Milk Room</u> - When used as a space contact spray for flies and other insects, remove animals from barn and close doors and windows tightly. Spray Purina Dairy Spray towards ceiling until a light fog hangs in air. Repeat as necessary.

<u>Beef Cattle, Hogs, and Other Farm Animals (Except Horses)</u> - Apply as a mist to the hair coat of farm animals for temporary repellent and killing action. Aids in controlling horn flies, house flies, mosquitoes and gnats. Do not wet the skin.

<u>CAUTION:</u> Harmful or fatal if swallowed. If swallowed do not induce vomiting. Call physician immediately. Avoid breathing of spray mist.

Do not use, pour, spill or store near an open flame. Do not wet the skin of animals. Avoid contamination of feed, foodstuffs, milk and milking untensils. Remove animals from barn when fogging or misting.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not use, pour, spill or store near an open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

EPA Est. No. 602-MO-1

QUALITY CONTROLLED BY PURINA RESEARCH

Printed in U.S.A.

9001 B