


PURINA®

E.P.A. Reg. No. 602-190

STOCK SPRAY-D

Contains Dioxathion

Kills... Horn Flies, Ticks, Lice, Sheep Keds



ions: ...
 other sprays ...
 dipping vat solutions ...
 dipping animals. Retain dipping vat ...
 ed. Clean out ...
 ined ...
 ticks. Do not spray horses or mules.
 ychroesterase ...
 f new thymus ...
 p. DMSO ...
 prn Flies, Lice, Ticks ...
 WARNING Do not dip
 Do not treat dairy animals
 Do not treat dairy goats
 Do not
 CATTLE BACK RUBBER
 HORN FLY CONTROL • AIDS IN LICE CONTROL
 Ections: Max 1 quart

ACCEPTED

JUL 7 1980

From the Federal Insecticide, Fungicide, and Rodenticide Act. Registered under EPA Reg. No. 602-190

ACTIVE INGREDIENTS	
Dioxathion 2,3,6-Dioxan-1,1-dithiolane-5-thiol-2-methylphosphorodithioate	11.9%
Related Compounds	5.0
Kyrene	50.0
Diethyl petroleum sulfonate	27.75
NEUT INGREDIENTS	5.35
	100.00%

KEEP OUT OF REACH OF CHILDREN
WARNING
 SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Manufactured by
RALSTON PURINA COMPANY

1000 ... 63186

STORAGE AND DISPOSAL
 INATE AND DESTROY USED CONTAINER

D-9023
 E.P.A. Est. No 602-MO-1

NET: 1 GAL.

45

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
WARNING

TREATMENT

ENVIRONMENTAL HAZARDS

PHYSICAL OR CHEMICAL HAZARDS

Mixing Directions

Use Directions

CATTLE - Horns

calves less than
SHEEP AND GOATS

HOGS - Lice T
treat sows within

STATEMENT OF PRACTICAL TREATMENT

DIRECTIONS FOR USE

SPRAY OR DIP

Water (1 gallon) Sprays 1 minute
Dip 10 minutes

Mixing Directions

Use Directions


pre-mix solution

CONTAMINATE

NOTE TO PHYSICIAN


WARNING - CONTAMINATE

QUALITY CONTROLLED BY PURINA RESEARCH



PURINA® STOCK SHIELD

Contains Dioxin
Kills... Horn Flies, Ticks, I



ACTIVE INGREDIENTS:
Dioxin 1.3 g. Thiocyanethion S.S.-bis
phosphorodithioate
Sulfated Castor Oils
Xylene
Aliphatic petroleum solvent
INERT INGREDIENTS

KEEP OUT OF REACH OF CHILDREN
WARNING
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

Manufactured by
RALSTON PURINA COMPANY
P.O. Box 107, The Keyboard Square

NET: 1 GALLON

Mixing Directions:

Use Directions:

CATTLE: Horn Flies, Lice, Ticks

calves less than 6 months of age. Do not treat dairy animals.

SHEEP AND GOATS: Lice, Ticks, Sheep Keds

HOGS: Lice, Ticks

treat sows with 12 weeks of farrowing or while nursing piglets.

CATTLE BACK RUBBER

FLY CONTROL • AIDS IN LICE CONTROL

Mixing Directions:

Use Directions:

STORAGE AND DISPOSAL

DO NOT CONTAMINATE ANIMALS OR FEEDS IN CONTAINER

D 9023

EPA Est. No. 602 MO 1

ENVIRONMENTAL HAZARDS:

PHYSICAL OR CHEMICAL HAZARDS:

DIRECTIONS FOR USE

SPRAY OR DIP

Lunge	Winter	Coast	Spouse	Other
Star	Tick	Tick	Ear	Tick
Tick				

ROLLED BY PURINA RESEARCH

ACCEPTED
 JUL 7 1980
 The Purina Company
 and Ralston Purina Co.
 Dept. of Agriculture
 and Food Inspection
 Service
 Washington, D.C. 20501