

ACCEPTED
 NOV 21 1972
 UNDER THE FEDERAL INSECTICIDE,
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PEST CONTROL
 REGISTERED UNDER NO. 501881
 TO ATTACHED COMMENTS.

SD-478 RALSTON PURINA CHEMICALS DIV 913 YELLOW RED BLACK

HOW TO USE PURINA WOUND PROTECTOR

Shake vigorously before using. Point valve opening toward wound in 17 oz. container 1 to 6 inches from wound while spraying.

To protect wounds: For castration, dehorning, docking, branding, wire or shear cut wounds, spray directly into the wound and 1 in. surrounding area. Treat navels of newborn animals. Apply to drainage area below wound of sheep and goats to prevent infestation. Re-treat at 5 to 7 day intervals until wounds are healed. When wounds become infected the supplemental use of a disinfectant solution may be necessary.

To kill and control screw worms, fleece worms and adult blow flies in and around superficial wounds: Spray wounds thoroughly with product material to penetrate into pockets made by maggots. Apply over the body discharge around wound to prevent re-infestation. Treat at 5 to 7 day intervals until wound is healed.

To kill spinose ear ticks: Spray directly into animal's ear. Repeat at 5 to 7 day intervals as needed.

CAUTION: Harmful or fatal if swallowed. Avoid contact with eyes, nose, mouth, or skin. Wash thoroughly after use. Avoid contact with skin.

Do not use on or around milk producing cats or other animals that lick themselves.

Keep out of reach of children. Do not use on or around children. Do not use on or around children. Do not use on or around children.


DANGER! Extremely flammable; contents under pressure. Do not use near fire, heated surfaces, sparks, or flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

729-1 D-9614 Printed in U.S.A.

EPA Reg. No. 602-191-AA

PURINA[®]
WOUND 
PROTECTOR

Protects Wounds Against Screwworms, Fleece Worms, Adult Blowflies, & Spinose Ear Ticks



WARNING: CONTENTS UNDER PRESSURE
CAUTION: KEEP OUT OF REACH OF CHILDREN
 READ CAREFULLY OTHER CAUTIONS ON BACK PANEL

Active Ingredients:
 Benzene (0.5% Ethyl 0.2% Trichlorophenyl) 2.80%
 Dipropyl succinate 4.00%
 Dibutyl sebacate 42.22%

Inert ingredients:
 Diethyl sebacate 50.00%



Date of Issue 2/19/45

NET WT. 1 OZ.

DISTRIBUTED BY RALSTON PURINA COMPANY
 St. Louis, Missouri 63103