Purina 602-209



18

1M-10

MAR 2 0 1980

Juder the Federal Insecticide, Implicide, and Rodentistic Aug or amended, for the posticide appropriate under PA Rea No. 2001

SPACE MIST INSECTICIDE

KILLS HOUSEFLIES, MOSQUITOES, GNATS, WASPS

Active Ingredients.

*(5-Benzyl-3-furyl) methyl 2,2-dimethyl-3-	
(2-methylpropenyl) cyclopropanecarboxylate	250%
Related compounds	.034%
Aromatic petroleum hydrocarbons	331%
Petroleum distillate	99 385%
	100.000%

*Cis-trans isomers ratio max 30% (±) cis and min 70% (±) trans

CAUTION

READ CAREFULLY OTHER CAUTIONS ON BACK PANEL

PURINA RESEARCH ASSURES QUALITY

RALSTON PURINA COMPANY

BEST AVAILABLE COPY

NET: 55 GALLON

017800 62394

PURINA' SPACE MIST INSECTICIDE

For use in food and beverage processing plants, food handling establishments, industrial and institutional establishments, hoppintals and nursing homes.

Harmful or fatal if swallowed. If swallowed, do not induce compting. Call physician immediately. Avoid breathing of spray mist. Doctor fremain in closed area being treated, ventilate area when treatment is completed.

Do not apply to uncovered human foodstuffs or food products. Avoid contamination of milk and milk equipment

ENVIRONMENTAL HAZARD

Cover fish bowls and remove pets while spraying

PHYSICAL OR CHEMICAL HAZARD

DO NOT USE, POUR. SPILL, OR STORE NEAR HEAT OR OPEN FLAME

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOUSEFLIES, MOSQUITOES, GNATS:

Close doors and windows to eliminate drafts. Direct mist from sprayer towards ceiling to contact flying insects. When using fixed misting or fogging devices, disperse Space Mist Insecticide horizontally below level of ceiling rafters. Use at the rate of 1/2 ounce for each 1,000 cubic feet of space. Keep building closed for 5 minutes after misting. Repeat spraying daily or as needed.

WASPS

Application should be made in late evening when insects are at rest Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

In hospitals and nursing homes, remove patients during application and ventilate rooms for two hours before reoccupying.

In processing plants, food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using Space Mist Insecticide as a space spray in these areas, apply only when plant is not in operation.

STORAGE AND DISPOSAL

Do not reuse empty container Destroy it by perforating or crushing. Bury or discard in a safe place.

LOT NO.