

# Purina®



Brand TM

EPA Reg. No. 602-209

Pg. 1 of 1  
Pm 1/7

# SPACE MIST INSECTICIDE

## KILLS HOUSEFLIES, MOSQUITOES, GNATS, WASPS

### ACTIVE INGREDIENTS

Resmethrin (5-Benzyl-3-furyl) methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate	250%
Related compounds	034%
Aromatic petroleum hydrocarbons	331%
Petroleum distillate	99.385%
	100.000%

10's trans isomers ratio: max. 30% (±) cis and min. 70% (±) trans

KEEP OUT OF REACH OF CHILDREN

### CAUTION

READ CAREFULLY OTHER CAUTIONS ON BACK PANEL

## PURINA® SPACE MIST INSECTICIDE

For use in food establishments and beverage processing plants, food handling establishments, industrial and institutional hospitals and nursing homes

### PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful or fatal if swallowed. If swallowed, do not induce vomiting. Call physician immediately. Avoid breathing of spray mist. Do not remain in closed area being treated. Ventilate area when treatment is completed.

Do not apply to uncovered human foodstuffs or food products. Avoid contamination of milk and milk equipment.

#### ENVIRONMENTAL HAZARD

Cover fish bowls and remove pets while spraying.

#### PHYSICAL OR CHEMICAL HAZARD

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**HOUSEFLIES, MOSQUITOES, GNATS:** Close doors and windows to eliminate drafts. Direct mist from sprayer towards ceiling to contact flying insects. When using fixed misting or fogging devices, disperse Space Mist Insecticide horizontally below level of ceiling rafters. Use at the rate of 1/2 ounce for each 1,000 cubic feet of space. Keep building closed for 5 minutes after misting. Repeat spraying daily or as needed.

**WASPS:** Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

In hospitals and nursing homes, remove patients during application and ventilate rooms for two hours before reoccupying.

In processing plants, food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using Space Mist Insecticide as a space spray in these areas, apply only when plant is not in operation.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

EPA Est. No. 602-MO-1

PURINA RESEARCH ASSURES QUALITY

4467 L

Printed in U.S.A.

# NET 55 GAL

LOT NO

**17800**

62394

