

HOW TO USE PURINA CAP-ZIN FUNGICIDE

Cap-Zin Fungicide is a combination of two preventive-type disease control chemicals that keep certain diseases from injuring roses, flowers, fruits, vegetables and lawns.

DIRECTIONS: Use 1½ tablespoons to 1 gallon of water (1 lb. to 50 gallons). Mix thoroughly and keep agitated while spraying.

FLOWERS <u>Roses</u> <u>Carnations</u> <u>Chrysanthemums</u>	<u>Black Spot</u> <u>Rust</u> <u>Leaf spot</u> <u>Flower blight</u> <u>Powdery mildew</u>	Begin applications before diseases appear or when new foliage appears. Spray upper and lower surfaces of all foliage thoroughly. Protect plants from early disease damage. Re-spray every 7 to 10 days and after rainy periods.
FRUITS <u>Apple</u> <u>Pear</u>	<u>Scab</u> <u>Frogeye</u> <u>Bitter rot</u> <u>Sooty blotch</u> <u>Fly speck</u>	Spray trees when flower buds begin to separate. Repeat at 7-10 day intervals until petal fall. Then repeat every 14 days. Do not spray within 7 days of harvest.
<u>Peach</u> <u>Plum</u>	<u>Scab</u> <u>Brown rot</u>	Apply when buds are full pink and keep blossoms protected. Respray when petals have dropped and at 10 day intervals. Do not spray within 30 days of harvest.
<u>Cherry</u>	<u>Leaf spot</u> <u>Brown rot</u>	Spray trees when most of the petals have fallen. Respray at 10 day intervals. Do not spray within 7 days of harvest.
<u>Raspberry</u> <u>Blackberry</u>	<u>Anthrachnose</u> <u>Spur blight</u>	Apply when blossoms are in bud (young canes 8 to 10 inches long). Respray 2 weeks later and in fall when old canes are removed. Do not apply within 14 days of harvest.
<u>Grape</u>	<u>Blackrot</u> <u>Downy mildew</u>	Apply just before bloom and immediately after (1 to 3 more sprays may be needed at 7 to 10 day intervals). Do not spray within 7 days of harvest.
VEGETABLES <u>Beans</u>	<u>Anthrachnose</u> <u>Rust</u> <u>Downy mildew</u>	
<u>Cucumber</u> <u>Melon</u> <u>Squash</u> <u>Pumpkin</u>	<u>Downy mildew</u> <u>Alternaria blight</u> <u>Anthrachnose</u> <u>Gummy stem blight</u>	Begin applications before diseases appear. Apply to upper and lower surfaces of all foliage. Protect plants from early disease damage. Repeat at 7 day intervals or as necessary. Do not apply within 7 days of harvest.
<u>Tomato</u> <u>Egg plant</u> <u>Pepper</u>	<u>Early blight</u> <u>Late blight</u> <u>Anthrachnose</u> <u>Septoria leaf spot</u>	
LAWNS TURF <u>Grass</u>	<u>Brown patch</u> <u>Damping-off</u> <u>Leaf spot</u>	Begin applications when new growth starts in spring. Apply 1 gallon per 100 sq. ft. of lawn area. Repeat at 7 to 10 day intervals as needed. Do not graze treated areas nor feed clippings to livestock.

WARNING: Avoid inhalation of dust or spray mist. May cause irritation of nose and throat. Avoid prolonged or repeated contact with skin. Do not contaminate food and feed. Do not allow mixture to stand. Apply only as specified on label. Burn empty bag. Keep out of smoke.

This product is toxic to fish. Keep out of any body of water.

QUALITY CONTROLLED BY PURINA RESEARCH

703D

G-3803

Printed in U.S.A.

TAKE TIME

OBSERVE LABEL
DIRECTIONS


602-211

602-211

ACCEPTED

JUL 20 1971

THE PURINA COMPANY
ST. LOUIS, MO. 63199



PURINA[®] CAP-ZIN[™] FUNGICIDE

FOR CONTROL OF PLANT DISEASES

ROSES • FLOWERS • FRUITS

VEGETABLES • LAWN



Active Ingredients:

*Captan	25.0%
Zineb (zinc ethylene bisdithiocarbamate)	25.0%
Inert Ingredients	50.0%
	100.0%

*N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS ON BACK PANEL

NET WT. 16 OZ. (1 LB.)

Manufactured by
RALSTON PURINA COMPANY
General Offices • Checkerboard Square • St. Louis, Mo. 63199