

602-213  
 RECEIVED  
 May 2 1974  
 DIVISION OF AGRI-CULTURE  
 REGISTERED

## HOW TO USE PURINA LIQUID FLY LARVICIDE MIXING DIRECTIONS FOR STRENGTH AND AMOUNT NEEDED

STRENGTH	1 GALLON	3 GALLONS	25 GALLONS	100 GALLONS
1/2%	3 oz.*	9 oz.	2 quarts	2 gallons
1%	6 oz.	18 oz.	1 gallon	4 gallons

Note: Agitate mixture thoroughly before application. \*2 Tablespoonfuls = 1 ounce.

### FLY CONTROL

MAGGOT CONTROL			FARM BUILDING SPRAY		
AREA	STRENGTH	APPLICATION	AREA	STRENGTH	APPLICATION
Caged Poultry, Manure Piles, and Other Fly Breeding Areas.	1%	Apply 1 gallon of mixture to every 100 square feet of droppings or surface area. Repeat as necessary.	Dairy, Poultry, Hog, Beef Cattle, Horse and Other Farm Buildings.	1%	Apply thoroughly to point of runoff to walls, ceilings and where flies congregate and feed. Use 1 gallon for each 500-1000 square feet. Repeat as necessary. Do not apply to birds or animals.

### POULTRY LICE AND MITE CONTROL

CAGED BIRDS			FLOOR BIRDS		
Wire Cages	1/2%	Apply directly 1 gallon of mixture to 100 birds using power sprayer at 100 psi. Or using low pressure sprayer, carefully apply 1 ounce of mixture per bird. Spray vent and fluff areas from below. Repeat when necessary but not more often than every 14 days.  Note: For maximum lasting control of the northern fowl mite, penetration of the feathers around the vent area is absolutely essential. Use power sprayer at the 100-125 psi at no less than recommended rate. More attention must be given each individual bird when using low-pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks. For positive control of mites on roosters dipping may be desirable.	Litter	1/2%	Apply 1 to 2 gallons per 1000 sq. ft. evenly to litter surface. Spray walls, roosts, cracks, crevices and interiors thoroughly. Spray birds lightly; treat roosters carefully to avoid reinfestation of breeding flocks.
			Roost Paint	1%	Use 1 pint to 100 feet. Apply thoroughly with brush or sprayer, especially to cracks.
			Fowl Tick (Blue Bug)	1%	Use 1 gallon for 100 to 150 sq. ft. Apply thoroughly to walls, ceilings, floor cracks and crevices with power sprayer.



### PRECAUTIONS IN USING

May be fatal if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.


**Note To Physician:** This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.



FOR N

**NEW  
FORMULA**

# PURINA® EPA Reg. No. 602-213-AA LIQUID FLY LARVICIDE




## PRECAUTIONS IN USING

May be fatal if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

**Note To Physician:** This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.



**Maggot Killer and Residual Fly Control. Also Controls Northern Fowl Mites, Chicken Mites and Lice On or Around Chickens**

### Active Ingredients:

2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	24.0%
Petroleum distillate	61.4%
Inert Ingredients	14.6%
	<u>100.0%</u>

Contains two pounds of active ingredient per gallon.

**CAUTION: KEEP OUT OF THE REACH OF CHILDREN  
SEE OTHER CAUTIONS ON BACK PANEL**

**NET CONTENTS: 1 GALLON**

**TAKE TIME**



**OBSERVE LABEL  
DIRECTIONS**

Printed in U.S.A.

MANUFACTURED BY  
**RALSTON PURINA COMPANY**  
GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63188