

EPA Reg. No. 602-214-AA



BOT CONTROL



NET WT. 0.72 oz. (FOR 0.18 oz. [5-Gram.] PACKETS)

SEE BACK OF BOX FOR COMPLETE INSTRUCTIONS AND WARNING STATEMENTS. KEEP OUT OF REACH OF CHILDREN.

Expiration Date Appears Inside on Each Packet.

4-PACK WORMS
ONE 1000 POUND
HORSE

ACTIVE INGREDIENT:
Dimethyl (2,2-
trichloro-1-hydroxy-
ethyl) phos-
phonate..... 90%
INERT INGREDIENTS:
(Colloidal Silica and
Dextrose)..... 10%
100%

DOSAGE. Give one packet (5 grams) per 250 pounds of body weight. For most effective control, repeat treatment at 3 to 4 month intervals. Purina BOT CONTROL FOR HORSES can be used at any time but do not repeat

NOTICE TO VETERINARIANS. Symptoms of overdosage in horses are ataxia and colic. Atropine is antidotal. Give 100 milligrams or 1½ grains atropine per 1,000 pounds body weight.

DOSAGE. Give one packet (5 grams) per 250 pounds of body weight. For most effective control, repeat treatment at 3 to 4 month intervals. Purina BOT CONTROL FOR HORSES can be used at any time but do not repeat more frequently than every 30 days.

DIRECTIONS. Use required dosage dry or dissolved in water. Mix with the amount of feed to be consumed at one feeding.

Withdrawal of feed 12 to 18 hours prior to treatment will assure complete ingestion. Allow horse to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours.

RESTRICTIONS. Use only as directed. Do not treat horses to be used for food. Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not exceed recommended dose. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using this product.

NOTICE TO VETERINARIANS. Symptoms of overdosage in horses are ataxia and Atropine is antidotal. Give 1/2 to 1 grain or 1 to 2 grains atropine per 1000 lbs. body weight.


WARNING. Not for human use. Keep out of reach of children. Harmful if swallowed. Avoid contact with the skin. Wash hands with soap and water after use. If swallowed, induce vomiting by giving 2 tablespoons of salt in a glass of water. Call physician immediately.

Purina Bot Control for Horses is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals.

TO PHYSICIAN. This material is a cholinesterase inhibitor. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Give supportive treatment. Do not walk patient nor give morphine.

NET WEIGHT: 0.72 oz. (FOUR 0.18 oz. [5-GRAM] PACKETS)

Expiration Date
and Lot No.
(On foil packet)



729G

F-0715

Distributed by

RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63188

Printed in U.S.A.

ACCEPTED

MAR 7 1973

THE FEDERAL INSECT

FU AND ODE CID CT

PO IN BT

LD TO IT LO HME S.

602-214

