

PURINA DISINFECTANT

Add 1 ounce (2 tablespoonfuls) Purina Disinfectant to each gallon of water. This solution contains more than 450 parts per million (ppm) of active ingredients. For bactericidal activity against Pseudomonas aeruginosa on porous surfaces and areas that are hard to clean, use 1-1/2 ounces per gallon of water. For Trichophyton interdigitale use 3 ounces per gallon of water.

USE DIRECTIONS

POULTRY AND LIVESTOCK BUILDINGS, DOG KENNELS, LABORATORIES, ZOOS: Remove all animals and feeds from premises, cars, trucks, coops, and crates. Remove all litter and droppings from floors, walls, and other surface areas of facilities. Add 1 ounce of Purina Disinfectant to each gallon of water and saturate all surface areas. Ventilate buildings and other closed areas before animals are housed. The disinfectant solution should be absorbed or dried.

HATCHERY OPERATIONS: Add 1 ounce of Purina Disinfectant to each gallon of water. After each hatch, follow cleaning with a liberal application of disinfectant solution. Spray the entire room after clean-up.

EGG SHELL SANITIZING: (Commercial and/or hatching eggs). To sanitize eggs which are not washed, use 2 ounces of Purina Disinfectant per 1 gallon of water (provides 920 ppm active quaternaries sanitizing solution). Apply with cloth, by spray, or by dipping. Rinse with potable water. Drain and air dry.

To sanitize eggs after washing, use 1½ ounces Purina Disinfectant per gallon of water. (provides 690 ppm active quaternaries sanitizing solution). Apply with cloth, by spraying, or by dipping. Rinse with potable water. Drain and air dry.

DAIRY SANITATION: Purina Disinfectant fulfills the criteria of Appendix F, the Grade A Pasteurized Milk Ordinance, 1965 recommendations of the U. S. Public Health Service in waters up to 750 ppm of hardness calculated as CaCO₃, when tested by the A.O.A.C. Germicidal and Detergent Sanitizers—Official Method.

Ask your local milk inspector or health officer for recom-

mendations on how to use Purina Disinfectant in his program.

Right after you milk, flush all equipment with cool or lukewarm water. Scrub equipment in a cleaning solution.

Then rinse equipment in a solution of Purina Disinfectant using 1 ounce (2 tablespoonfuls) to each 2¼ gallons of water. Soak at least 2 minutes, allow equipment to dry, and protect from contamination. *At this level, no potable water rinse is required.*

SWIMMING POOLS: As a disinfectant for swimming pool surfaces (walks and runways of supporting surfaces, use 1½ ounce Purina Disinfectant per gallon of water (provides 690 ppm active quaternaries). Apply with mop, by spraying, or by flooding surface. Allow to air dry.

ALGAEICIDE FOR SWIMMING POOLS: Starting with a clean freshly filled pool, add 1 tablespoon of Purina Disinfectant to each 100 gallons of water. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae is still present or reappears, repeat treatment procedure as necessary. When pool is free of visible algae, add a maintenance dose of 1 tablespoonful per 400 gallons of water.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail, or pool has unusually heavy use, add treatment more frequently.

HOME DISINFECTION: Add 1 ounce of Purina Disinfectant per gallon of water to disinfect and deodorize refrigerators, freezers, garbage pails, bathrooms, and floors.

NOTE: Do not use with soap or other cleaners.

WARNING: Causes eye and skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Rinse empty container thoroughly with water and discard.

PROTECT FROM FREEZING

EPA Est. No. 602-MO-1

A-2804

ACCEPTED

FEB 25 1976

Under the Federal Insecticide,

Fungicide, and Plant Growth Regulator Act.

recommendations on how to use Purina Disinfectant in his program.

Right after you milk, flush all equipment with cool or lukewarm water. Scrub equipment in a cleaning solution.

Then rinse equipment in a solution of Purina Disinfectant using 1 ounce (2 tablespoonfuls) to each 2 1/4 gallons of water. Soak at least 2 minutes, allow equipment to dry, and protect from contamination. *At this level, no potable water rinse is required.*

SWIMMING POOLS: As a disinfectant for swimming pool surfaces (walks and runways of supporting surfaces, use 1 1/2 ounce Purina Disinfectant per gallon of water provides 690 ppm active quaternaries). Apply with mop, by spraying, or by flooding surface. Allow to air dry.

ALGAEICIDE FOR SWIMMING POOLS: Starting with a clean freshly filled pool, add 1 tablespoon of Purina Disinfectant to each 100 gallons of water. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae is still present or reappears, repeat treatment procedure as necessary. When pool is free of visible algae, add a maintenance dose of 1 tablespoonful per 400 gallons of water.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail, or pool has unusually heavy use, add treatment more frequently.

HOME DISINFECTION: Add 1 ounce of Purina Disinfectant per gallon of water to disinfect and deodorize refrigerators, freezers, garbage pails, bathrooms, and floors.

NOTE: Do not use with soap or other cleaners.

WARNING: Causes eye and skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Rinse empty container thoroughly with water and discard.


PROTECT FROM FREEZING

EPA Est. No 602-MO-1

A-2804

BY PURINA RESEARCH

Printed in U.S.A.




TM

PURINA

DISINFECTANT

**A General Purpose Disinfectant for Farm and
Livestock and Poultry Buildings and Equipment • Hotel
Home Sanitation • Milking Machines and Dairy Utensils
Zoos, Dog Kennels, and Fur Bearing Operations • Eggs**



Active Ingredients:

n-Alkyl (60% C₁₄, 80% C₁₆, 5% C₁₂, 5% C₁₀)

dimethyl benzyl ammonium chlorides.....

n-Alkyl (68% C₁₂, 82% C₁₄) dimethyl ethylbenzyl

ammonium chlorides.....

Inert Ingredients.....

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL WARNINGS**

NET CONTENTS: 32 FL. OZ.

MANUFACTURED BY

RALSTON PURINA CO.
GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MO.