


602-221

10/24/1972

23




easy to apply

E.P.A. REG. NO.
602-221-AA

PURINA® CATTLE DUST

DAIRY CATTLE — BEEF CATTLE



Active Ingredient:

*Methoxychlor, Technical.....	10.0%
Inert Ingredients.....	90.0%
	<u>100.0%</u>

*Equivalent to 8.8% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 1.2% other isomers and related compounds.



CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON BACK PANEL

NET WT. 20 OZ. (1¼ LB.)

Manufactured By
RALSTON PURINA COMPANY
General Offices • Checkerboard Square • St. Louis, Missouri 63188

**HOW TO USE
PURINA CATTLE DUST**

Dust livestock in the spring when hornflies first appear to prevent rapid increase in number of flies.

DIRECTIONS:

Milk Cows—Sprinkle Cattle Dust over the shoulders and back. Rub dust lightly into the haircoat. Use one pound to 10 animals. Treat every 3 weeks, if necessary, for best hornfly control.

Calves and Heifers—Sprinkle Cattle Dust over shoulders, back and sides. Rub lightly into the haircoat. Use one pound to 15-30 animals, depending on size. Treat as often as necessary for best hornfly control.

Beef Cattle—Sprinkle Cattle Dust over shoulders, back and sides. Work thorough application into the hair coat. Treat as often as necessary for best hornfly control.

CAUTION: Avoid contamination of feed and foodstuffs. Protect milk and milking equipment from contamination. Do not inhale.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Empty bags should be buried or burned. If burned, stay out of smoke and bury ashes in a safe place.

77474
10-10-72

ACCEPTED
OCT 24 1972
UM
FUNGICIDE AND RIDOMIL IDE ACT
FOR ECONOMIC PESTICIDE REG.
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

602-221