NEW FORMULA

E.P.A. REG. NO.
602-227-AA

PURINA HOG & CATTLE DUSTING POWDER

CONTROLS LICE, HORNFLIES & TICKS ON PIGS • DAIRY OR BEEF CATTLE

Active Ingredient:	
*Malathion	4.0%
Inert Ingredients	96.0%
	100.0%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON BACK PANEL

NET WT. 16 OZ. (1 LB.)

TAKE TIME
OBSERVE LABEL DIRECTIONS

MANUFACTURED BY
RALSTON PURINA COMPANY
GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63188

HOW TO USE PURINA HOG AND CATTLE DUSTING POWDER

SWINE

HOG LICE: For the control of lice on swine.

Thoroughly apply Hog and Cattle Dusting Powder to all animals of the herd. In addition, pens should be thoroughly dusted.

For pigs less than one month of age: Apply ¼ to ½ tablespoonful.

One treatment may be sufficient. Repeat only as needed.

DAIRY OR BEEF CATTLE

HORN FLIES--Apply 4 tablespoonfuls of Hog and Cattle Dusting Powder on the back and neck of the animal. Repeat applications at 10 to 14 day intervals.

Do not apply to dairy cattle later than 5 hours before milking or during milking.

NON-LACTATING DAIRY COWS OR BEEF CATTLE

LICE, TICKS--Dust animal thoroughly. Repeat treatment after 2 to 3 weeks if needed.

Do not apply thorough dusting for lice or ticks on lactating dairy animals, or non-lactating dairy cows within 2 weeks of freshening.

Dust calves lightly.

CAUTION: Avoid breathing of dust. Avoid contact with skin, eyes or clothing. Do not leave dust on skin; do not get dust in eyes. Wash thoroughly after using. Keep out of reach of children.

Avoid contamination of feed, food containers and watering troughs.

This product is toxic to fish and birds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

D-4052 Printed in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH