

Front Panel

BEST AVAILABLE COPY

PURINA [®]
BRAND

HOG AND CATTLE
DUSTING POWDER

EPA Reg. No. 602-227

(3-row Checkerband) [®]

CONTROLS LICE, HORNFLIES & TICKS

ON

PIGS, DAIRY OR BEEF CATTLE

ACTIVE INGREDIENT:

*Malathion 4.0%

INERT INGREDIENTS 96.0%

100.0%

*O-O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Back Panel For Precautionary Statements

NET WT 16 OZ (1 LB)

D-4052


(Note: Type size and prominence will conform with requirements for precautionary labeling.)

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 10 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

602-227



NET 16 OZ
PURINA [®] HOG & CATTLE
BRAND
DUSTING POWDER

Manufactured By
RALSTON PURINA COMPANY
General Offices
Checkerboard Square
St. Louis, Missouri 63164

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

BEST AVAILABLE COPY

CAUTION

Avoid breathing dust or spray mist. Avoid contact with skin or clothing. Wash skin thoroughly after handling. Remove and wash contaminated clothing before reuse.

In case of contact, wash skin with plenty of soap and water; For eyes, flush with water for 15 minutes. Get medical attention if irritation persists.

Note to Physician: Malathion is an acetyl cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SWINE

HOG LICE: For the control of lice on swine.

Thoroughly apply Hog and Cattle Dusting Powder to all animals of the herd. In addition, pens should be thoroughly dusted. One treatment may be sufficient. Repeat only as needed. Do not treat animals under one month of age. Do not contaminate feed, food containers and water/or water troughs during application.

DAIRY OR BEEF CATTLE

HORN FLIES - Apply 4 tablespoonfuls of Hog and Cattle Dusting Powder on the back and neck of the animal. Repeat applications at 10 to 14 day intervals. Dust calves lightly. Do not treat calves under one month of age. Do not contaminate feed, food containers and water/or water troughs during applications.

Do not apply to lactating dairy cattle later than 5 hours before milking. Ideal time for application is immediately after milking.

NON-LACTATING DAIRY COWS OR BEEF CATTLE

LICE, TICKS - Dust animal thoroughly. Repeat treatment after 2 to 3 weeks if needed. Dust calves lightly. Do not treat calves under one month of age. Do not contaminate feed, food containers and water/or water troughs during applications.

Do not apply to non-lactating dairy cattle within 2 weeks of freshening.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

QUALITY CONTROLLED BY PURINA RESEARCH

EPA Est. No. 602-MO-1

8401 E

Printed in U.S.A.

BEST AVAILABLE COPY