

HOW TO USE PURINA FLY BAIT GRANULES

Scatter Purina Fly Bait Granules daily or as needed to reduce fly population. Distribute bait from container or with jar or can to avoid handling bait. Keep Fly Bait Granules in original container tightly closed and store in cool place.

Loafing Sheds, Livestock Pens, Pig Parlors, Stables, Broiler Houses: Use only around outside areas. Scatter Fly Bait Granules at the rate of $\frac{1}{4}$ lb. per 1000 sq. ft. on ground areas where flies congregate. Ideal baiting sites are around doorways, along fence lines, soiled areas, or in alleyways of feedlots.

Caged Poultry and Turkeys on Wire: Scatter Fly Bait Granules on walkways at the rate of $\frac{1}{4}$ lb. per 500 sq. ft. to aid in controlling house flies. Also scatter outdoors around doorways.

Dog Kennels and Furbearing Animal Units: Scatter Fly Bait Granules around outside areas of dog pens, runways, and other surfaces out of reach of the animals. For cage units, scatter on walkways at the rate of $\frac{1}{4}$ lb. per 500 sq. ft.

Food, Poultry and Meat Processing Plants: Scatter Fly Bait Granules at the rate of $\frac{1}{4}$ lb. per 1000 sq. ft. in inedible departments only. Fly Bait Granules may be used outside livestock pens or loading and unloading docks.

WARNING:

May be harmful or fatal if swallowed, inhaled or absorbed through skin. **DO NOT** get on skin,

in eyes, or on clothing. Wash thoroughly with soap and water after using. Keep out of reach of children.

Do not contaminate feed and foodstuffs. Do not apply where poultry or animals, especially young calves, can pick it up or lick it. Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Use only in areas *inaccessible* to food producing animals.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Use this product only as specified on this label.


Do not re-use empty container. Destroy container by perforating and bury or discard in a safe place.

Should illness occur, get prompt medical aid.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal.



ANTIDOTE: If swallowed induce vomiting with 1 tablespoonful of salt and warm water.

QUALITY CONTROLLED BY PURINA RESEARCH



PURINA®
FLY BAIT
GRANULES
ATTRACTS and KILLS HOUSE FLIES

EPA REG. NO.
 602-229-AA



Active Ingredient:

*Dimethyl 3-hydroxyglutaconate dimethyl phosphate†.....	1.0%
Inert Ingredients.....	99.0%
	<u>100.0%</u>

*Bomyl *Reg. TMAllied Chemical Corporation †U.S. Patent No. 2,891,887

KEEP CONTAINER CLOSED • STORE IN COOL DRY PLACE

WARNING: KEEP OUT OF REACH OF CHILDREN
 SEE OTHER WARNINGS ON BACK PANEL



NET WT.: 48 OZ. (3 LBS.)

Manufactured by
RALSTON PURINA COMPANY
 GENERAL OFFICES • CHECKERBOARD SQUARE • ST. LOUIS, MISSOURI 63188