


662-231



EPA Reg. N
40231 AA

PURINA® ROSE AND FLOWER DUST

CONTROLS INSECTS AND PLANT DISEASES



CAUTION: ...

NET WT. 10 OZ.

RASTON PURINA COMPANY

PURINA ROSE AND FLOWER DUST

DIRECTIONS

Apply Purina Rose and Flower Dust to the upper and lower leaf surfaces of foliage for complete coverage. Protect plants from early insect and disease damage. Begin application before diseases appear. Repeat applications at 7 to 14 day intervals for disease control; repeat as necessary for insect control. Dust when air is calm. For use in flowers and ornamental plants only.

CONTROLS: Insects—Aphid, Fuller rose beetle, rose chaffer, rose budworm, Todd callosity, rose slug, flea beetle, Japanese beetle, lace bug, leafhopper, leaf miner, plant hopper, spider mite, bagworms, scale crawlers, spotted leaf beetles, white flies.

Plant Diseases:

Powdery Mildew—after the plants have been cleaned, saturated with water and allowed to dry. Use on snapdragons, zinnia.

Black Spot—rose.

Anthraxnose—hollyhock, geranium.

Botrytis Blight—gallium, geranium.

Leaf Spot—geranium, geranium.

Rust—hollyhock, geranium.

Leaf Blight—zinnia.

NOTE: Injury to tender tips may occur if plants are wet when dusted. The presence of high humidity, wind, or frost may reduce the effectiveness of Japanese beetle. The product may be toxic to bees, and should not be applied to flowers in bloom. For more information, write to Purina Research, P.O. Box 1356, St. Louis, Mo. 63103.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of dust. Avoid contact with skin and eyes. If contact occurs, wash thoroughly. Wash thoroughly after handling. Keep out of reach of children and domestic animals. Avoid contact with food and feed.

Note for Physician: ...

ANTIDOTE: ...

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

G 89-01

QUALITY CONTROLLED BY PURINA RESEARCH



Active Ingredients

*Carbaryl (1-naphthyl N-methylcarbamate).....	3.000%
*Malathion.....	4.000%
Zinc ion and manganese ethylene bisdithiocarbamate.....	4.000%
(A coordination product of Manganese 0.8%, Zinc 0.1%, and ethylene bisdithiocarbamate 3.1%)	
2,4-Dinitro-6-octylphenyl crotonate.....	0.812%
2,4-Dinitro-4-octylphenyl crotonate.....	0.812%
Nitrooctyl phenols (principally dinitro).....	0.056%
Inert Ingredients.....	88.132%
	100.000%

*SEVIN Registered Trademark of Union Carbide Corporation U.S. Pat. No. 2,915,178

**Dithiocarbamate salts of dimethylmercaptoacetate

***A mixture of 2-n-octylheptyl, 1-ethylhexyl, and 1-propylheptyl esters. KARATHANE Registered Trademark of Ralston Purina Company U.S. Pat. No. 2,570,660

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON BACK PANEL**

NET WT.: 10 OZ.

Manufactured by

RALSTON PURINA COMPANY

General Offices: 500 North Lincoln Avenue • St. Louis, Missouri 63188

Leaf Blight—zi

NOTE: Injury to wet when dusted. Do not dust Box Maple. This product for direct treatment. For more information, see Agricultural Extension

CAUTION: Harmful if inhaled. Avoid breathing of dust. May be absorbed through skin. Avoid handling. Keep away from children and animals. Avoid

Note for Phytotherapy: Moderate, repeated use. Atropine is an antidote.

ANTIDOTE: If inhaled, give in a glass of water and water; for eye contact, get medical attention.

This product is for use on trees and shrubs. Do not use on lawns or ponds. Do not contaminate water. Do not use on or disposal of this product as specified on the label. Do not reuse empty containers. Do not bury with waste. **KEEP OUT OF REACH OF CHILDREN**

QUALITY CONTROL