

PM15-602-240

Prev. Date Issued: 2/3/88

10/11

PURINA
BRAND

DIAZINON*INSECTICIDE-25E

RECOMMENDED FOR AGRICULTURAL/COMMERCIAL USE

(Must be printed at top of front panel)

EPA Reg. No. 602-240
EPA EST. No. 602-MO-1

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.

(3-Row Checkerband [®])

PROTECTS
ROSES-FLOWERS-TREES-SHRUBS-
LAWNS-VEGETABLES-FRUITS-NUTS
CONTROLS
NUISANCE PESTS LISTED ON INSERT IN OUTSIDE AREAS
AND AS A PREMISE FLY SPRAY

ACTIVE INGREDIENT:

Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	25.0%
INERT INGREDIENTS	75.0%
	100.0%

KEEP OUT OF REACH OF CHILDREN
WARNING
SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON BACK PANEL

PRECAUCION

Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicaca ampliamente.

(English Translation)

CAUTION

To the User: If you cannot read English, do not use this product until the label has been fully explained to you.

NET 32 FL OZ (1 QT)

G-2657


(9-Square Logo)

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812
St. Louis, Missouri 63166-6812 U.S.A.

NET 32 FL OZ

PURINA [®] DIAZINON*
BRAND
INSECTICIDE-25E

ACCEPTED
SEP - 2 1988
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 602-240



10/11

Back Panel

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
WARNING

Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals. DO NOT USE IN THE HOME. Do not contaminate ornamental fish ponds.

Do not apply this product in such a manner as to directly or through drift expose workers of other persons. The area being treated must be vacated by unprotected persons.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed - Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin - Wash immediately with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately.

Note to Physician - Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally. If symptoms of cholinesterase inhibition are present, atropine sulfate is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

ENVIRONMENTAL HAZARD

This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

STORAGE AND DISPOSAL

Storage: Store unused PURINA Diazinon Insecticide-25E in original container only, in cool, dry area out of reach of children and animals.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.


Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

NOTE: See Directions For Use on Insert

QUALITY CONTROLLED BY PURINA RESEARCH

8802 P

Printed in U.S.A.



3 of 11

Reason For Issue: For EPA approval

Date Issued: 6/17/88 (White)

Prev. Date Issued: 2/3/88

4

Front Copy Of Insert For Pt. & Qt.

(9-Square Logo)

PURINA[®]
BRAND

DIAZINON*INSECTICIDE-25E

RECOMMENDED FOR AGRICULTURAL/COMMERCIAL USE

EPA Reg. No. 602-240

THIS PRODUCT MUST NOT BE USED ON GOLF
COURSES AND SOD FARMS.

EPA EST. No. 602-MO-1

PROTECTS
ROSES-FLOWERS-TREES-SHRUBS-
LAWNS-VEGETABLES-FRUITS-NUTS
CONTROLS
NUISANCE PESTS IN OUTSIDE AREAS
AND AS A PREMISE FLY SPRAY

KEEP OUT OF REACH OF CHILDREN
WARNING
SEE LABEL FOR PRECAUTIONARY STATEMENTS

ACCEPTED
SEP - 2 1988
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 602-240

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812
St. Louis, Missouri 63166-6812 U.S.A.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

40/11

Inside Copy of Insert**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Directions for Mixing and Applying PURINA[®] DIAZINON^{*} Insecticide-25E

Mix and spray as directed in rate chart. Use a coarse drenching spray for lawns, soil and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Repeat application as needed or as noted in the rate chart. Flush sprayer with clean water after each use. Use contents within 6 months after opening.

Problem Areas**Pests - Rates and Directions**

Note: Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia and orchids since injury may occur.

ROSES AND FLOWERS

Azaleas, Camellias,
Chrysanthemums, Carnations,
Gladioli, Lilacs,
Roses, Rhododendrons

Aphids, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Dipterous Leafminers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Twospotted Spider) Thrips, Whiteflies

1 fl. oz. (2 tablespoons) per 3 gals. of water.

Apply as a thorough foliar spray. Wet the plant to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

TREES AND SHRUBS

Arborvitae, Azalea, Birch,
Boxwood, Camellia
Douglas Fir, Elm, Hawthorn,
Holly, Juniper, Lilac,
Locust, Maple, Mimosa, Oak,
Pine, Plum, Poplar,
Rhododendron, Spruce,
Willow

Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak) Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Twospotted Spider), Obscure Root Weevils, Omnivorous Leaf-tiers, Scale Crawlers (Cottony-cushion, Euonymus, Lancaium, Pine Needle, Pit Scale, San Jose, Soft), Webworms (Juniper, Mimosa *), Whiteflies.

1 fl. oz. (2 tablespoons) per 3 gals. of water.

Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

***Mimosa Webworms:** Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.

Apple-and-Thorn Skeltonizers, Hemlock Chermes, Oak Loopers, Oblique-handed Leaf Rollers, Pear Slugs, Tent Caterpillars, Webworms (Cotoneaster, Fall)

3 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray.

LAWNS

Bent, Bermuda, Bluegrass,
Dichondra, Fescue,
Merion, St. Augustine

Fire Ants

To aid in the control of fire ants in lawns and other recreational areas, apply 1 fl. oz. (2 tablespoons) in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. **Note:** For best results, apply in cool weather 65-80°F or in early morning or late

Inside Copy of Insert - ContinuedProblem AreasPests - Rates and DirectionsLAWNS (Cont.)

Ants, Armyworms, Bermudagrass Thrips, Brown Dog Ticks, Chiggers, Crickets, Cutworms, Earwigs, Fleas, Fruit Flies, Lawn Billbugs, Chinch Bugs, Lawn Moths (Sod Webworms), Leafhoppers, Mites (Bermuda, Clover), Sowbugs (Pillbugs), Springtails (Collembola)

1 fl. oz. in 3 gals. of water covers 125 sq. ft.

8 fl. oz. in 24 gals. of water covers 1,000 sq. ft.

Thoroughly wet down grass a few hours before applying.

⁺ Brown Dog Ticks: Do not apply to animals. In particular, spray under shrubbery and the grass next to house.

⁺⁺ Lawn Billbugs: Water the lawn thoroughly immediately after treatment.

⁺⁺⁺ In the North, if Chinch Bugs are the only lawn insect problem, a lower rate of application can be used: 1/2 fl. oz. in 3 gals. of water covers 125 sq. ft.

Millipedes: 2 fl. oz. in 3 gals. of water covers 125 sq. ft.

Thoroughly wet grass a few hours before applying.

Rhodesgrass Scales

1 fl. oz. in 3 gals. of water covers 100 sq. ft.

Apply when Scale Crawlers first emerge.

Ants, Digger Wasps

1 fl. oz. in 3 gals. of water covers 125 sq. ft.

Spot spray ant hills and wasp ground nest openings.

⁺ Make the application when the wasps are not active.

Billbugs⁺⁺, White Grubs⁺⁺⁺, of: Japanese Beetle, European Chafer, Southern Chafer

1 fl. oz. in 3 gals. of water covers 125 sq. ft.

Water grass thoroughly after application.

⁺⁺ Billbug control: Apply when activity is first observed or when chewed or brown grass is present, indicating damage from the insect.

⁺⁺⁺ Grub control: Make application any time between late July and early October.

Hyperodes Weevil (New York only)

3/4 fl. oz. in 3 gals. of water covers 125 sq. ft.

Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

6/11

Inside Copy of Insert - Continued

1

Problem Areas	Pests - Rates and Directions
Vegetables* Foliar Application:	Thoroughly spray all plant parts. Generally, applications at 7 to 10-day intervals may be required. Do not feed treated vines to livestock.
Beans, Peas	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites. 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Cabbage, Cauliflower, Collards, Broccoli, Kale, Radishes, Turnips	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage Worms, Dipterous Leaf Miners 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Carrots, Celery, Endive, Lettuce, Parsnip, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leaf Miners, Mites 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Cantaloupes, Cucumbers, Muskmelon, Squash (summer, winter), Watermelon	Aphids, Thrips, Mites 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Peppers, Potatoes, Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms (tomato), Flea Beetles, Leafhoppers, Southern Armyworms (potato and tomato), Vinegar Flies (Drosophila spp. [tomato]) 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Onions	Onion Thrips 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Sweet Corn	Aphids, Flea Beetles, Grasshoppers, Mites 1 fl. oz. in 3 gals. of water covers approximately 1,000 sq. ft. of garden.
Vegetables* Preplant Soil Application:	Cutworms (Surface, Subterranean) 6 fl. oz. per 1,000 sq. ft. in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 2-3 inches for Surface Cutworms, 3-6 inches for Subterranean Cutworms. Garden Symphylans (Garden Centipedes) 7-1/2 fl. oz. per 500 sq. ft. in sufficient water to achieve thorough coverage. Apply broadcast just before planting. Immediately work into soil 4-8 inches.

7/11

Inside Copy of Insert - Continued

8

Problem Areas	Pests - Rates and Directions
Vegetables* Preplant Soil Application: (Continued)	Mole Crickets: Carrots, Corn, Peas, Pole Beans, Red Beets, and Snap Beans <u>Only</u> .
	1-1/2 fl. oz. per 1,000 sq. ft. in sufficient water to achieve thorough coverage
	Apply broadcast just before planting. Apply 1-2 days before sowing or transplanting. Immediately work into soil 1-2 inches.
	Root Maggots: Cabbage <u>Only</u>
	4-1/2 fl. oz. per 1,000 sq. ft. in sufficient water to achieve thorough coverage.
	Apply broadcast just before planting. Immediately work into soil 3-4 inches before seeding, either in seed bed or in field or before transplanting.
	Wireworms: Cabbage, Carrots, Corn, Lettuce, Peas, Snap Beans, & Tomatoes <u>Only</u>
	6 fl. oz. per 1,000 sq. ft. in sufficient water to achieve thorough coverage.
	Apply broadcast just before planting. Immediately work into soil 4-8 inches.
	*After spraying, do not pick vegetables within:
	1 day - Peas, Tomatoes, Sweet Corn
	3 days - Cantaloupes, Muskmelons, Winter Squash, Watermelons
	5 days - Broccoli, Cauliflower, Peppers
	7 days - Beans, Cabbage, Cucumbers, Summer Squash
	10 days - Carrots, Celery, Collards, Endive, Kale, Lettuce, Onions, Parsnips, Radishes, Spinach, Turnips
	35 days - Potatoes
Fruits⁺ & Nuts	Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted. If infestations warrant, repeat applications at 7 to 10 day intervals.
Almonds, Filberts, Pecans, Walnuts	Aphids, Nut Casemakers (pecans) Scale Crawlers, Spider Mites
	1/4 fl. oz. in 3 gals. of water. Apply as a full cover spray.
	Do not apply after husks open or split.

80/11

9

Inside Copy of Insert - Continued

Problem Areas	Pests - Rates and Directions
Apples, Pears	Aphids, Codling Moths, Eye-spotted Bud Moths, Fruit Tree Leaf Rollers, Leafhoppers, Tentiform Leaf Miners (pears), Maggots (apples), Mealybugs, Mites (European Red, Clover [brown], Pacific, Two-spotted Spider), Psylla (pears), Scale Crawlers (Cottony-Cushion, San Jose, Forbes, Soft) 1 fl. oz. in 3 gals. of water. Apply as a full cover spray. On apples, slight russeting may occur on some varieties such as Golden Delicious.
Apricots, Peaches,	Aphids, Apricot Mealybugs (apricot), Mites (clover [brown], Two-spotted Spider), Scale Crawlers (Olive, San Jose) 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Cherries	Black Cherry Aphids, Cherry Fruit Flies, Cherry Fruitworms (Maggots), Fruit Tree Leaf Rollers, Mites (Cherry Rust), Scale Crawlers (San Jose) 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Figs	Dried Fruit Beetles, Vinegar Flies (<i>Drosophila</i> spp.) 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Plums, Prunes	Aphids (Leaf Curl, Plum, Mealy Plum, Thistle), Mites (Clover [brown], European Red) 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Caneberries (blackberries, boysenberries, dewberries, loganberries, raspberries)	Aphids, Leafhoppers, Raspberry Sawflies, Raspberry Fruitworms, Two-spotted Mites, Thrips 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Blueberries	Aphids, Blueberry Maggots, Cranberry Fruitworms, Cherry Fruitworms, Two-spotted Mites, Thrips 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Strawberries	Aphids, Two-spotted Mites, Strawberry Leaf Rollers 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Grapes	Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers, Scale Crawlers 1 fl. oz. in 3 gals. of water. Apply as a full cover spray.

9 of 11

Inside Copy of Insert - Continued

<u>Problem Areas</u>	<u>Pests - Rates and Directions</u>
Grapefruit, Lemons, Oranges	<p>Citrus Thrips, Scale Crawlers (Citrus Snow⁺⁺, Cottony-cushion)</p> <p>1 fl. oz. in 3 gals. of water. Apply as a full cover spray.</p> <p>⁺⁺ 1 fl. oz. in 3 gals. of water. Apply as a full cover spray after blooming and again in the summer. Thorough spray coverage of tree trunk, limbs and twigs is essential.</p> <hr/> <p>⁺ After spraying do not pick fruit within:</p> <p>5 days - Strawberries, Figs 7 days - Blueberries, Caneberries 10 days - Apricots, Cherries, Grapes, Nectarines, plums, Prunes 14 days - Apples, Pears 21 days - Grapefruit, Lemons, Oranges, Peaches</p> <hr/> <p>Nuisance Pests in Outside Areas</p> <p>Ants⁺, Boxelder Bugs, Brown Dog Ticks⁺, Chiggers⁺, Clover Mites⁺, Cockroaches⁺, Crickets⁺, Earwigs⁺, Fleas⁺, Flies⁺, Millipedes⁺, Sowbugs (Pillbugs)⁺, Spiders, Springtails (Collembola)⁺</p> <p>8 fl. oz. in 3 gals. of water.</p> <p>Thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entranceways, walks, garbage cans, tree trunks and into cracks and other places where insects can hide.</p> <p>Do not treat plants or animals.</p> <p>* Spray a 5 ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. See lawn recommendations for other control measures for these pests.</p>

Recommended for Agricultural/Commercial Use

Fly control in barns and animal sleeping quarters (except dairy barns, milk rooms, and poultry houses).

Residual Sprays: Remove animals from buildings prior to treatment and keep animals out for at least 4 hours. Apply as an overall spray to ceilings and walls of livestock sheds, calf barns, hog barns, loafing sheds, and other farm buildings except poultry houses, dairy barns and milk rooms. Areas where flies congregata, such as garbage dumps and corrals, should be sprayed thoroughly. Applications should be made to point of run off. One gallon of the above spray should cover from 350-750 ft. of area, depending upon the surface treated. Repeat applications as fly populations warrant.

