

IODINE CONCENTRATE 3.5

P-0114

DIRECTIONS:

1. For general disinfection of walls, floors, and woodwork in public buildings, institutions, and hospitals. Thoroughly clean surfaces to be disinfected. Use a solution of 1 ounce of PURINA IODINE CONCENTRATE 3.5 per 4 gallons of water (1:512). Apply disinfecting solution with a clean mop or cloth so as to wet surfaces thoroughly. If Tuberculocidal activity is desired, use a solution of 1 ounce of PURINA IODINE CONCENTRATE 3.5 per 2 gallons of water, and follow procedures above.


2. For sanitization of equipment in the food processing industry, thoroughly clean all surfaces to be sanitized. Use a solution of 1 ounce of PURINA IODINE CONCENTRATE 3.5 per 12 gallons of water (1:1536). Apply sanitizing solution with a clean mop, cloth, or through power equipment so as to wet surface thoroughly. Rinse with potable water.

WARNING: Keep Out Of Reach Of Children. Causes eye or skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

Rinse empty container thoroughly with water and discard.



Purina

IODINE CONCENTRATE 3.5

P-0114

DIRECTIONS:

1. For general disinfection of walls, floors, and woodwork in public buildings, institutions, and hospitals. Thoroughly clean surfaces to be disinfected. Use a solution of 1 ounce of PURINA IODINE CONCENTRATE 3.5 per 4 gallons of water (1:512). Apply disinfecting solution with a clean mop or cloth so as to wet surfaces thoroughly. If Tuberculocidal activity is desired, use a solution of 1 ounce of PURINA IODINE CONCENTRATE 3.5 per 2 gallons of water, and follow procedures above.

2. For sanitization of equipment in the food processing industry, thoroughly clean all surfaces to be sanitized. Use a solution of 1 ounce of PURINA IODINE CONCENTRATE 3.5 per 12 gallons of water (1:1536). Apply sanitizing solution with a clean mop, cloth, or through power equipment so as to wet surface thoroughly. Rinse with potable water.

WARNING: Keep Out Of Reach Of Children. Causes eye or skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

Rinse empty container thoroughly with water and discard.


EPA Reg. No. 602-244-AA

P-0114

IODINE CONCENTRATE 3.5

602-244

**WARNING: KEEP OUT OF REACH OF CHILDREN
CAUSES SKIN AND EYE IRRITATION
SEE OTHER WARNINGS ON SIDE PANEL**



Active Ingredients:

Alpha (p-nonylphenyl)-omega-hydroxypoly-oxoethylene-iodine complex providing 3.5% titratable iodine	27.60%
Phosphoric Acid	9.00%

Inert Ingredients:

63.40%

U.S. Patent Nos. 2,941,777-3 078,799

**CHEMICAL
SPECIALTIES™**

NET CONTENTS: 1 GALLON

MADE IN U.S.A.

PU
ver
bac
Sci
Th
tiob
Sta
Sal
Pac
My
Tri
Sta
Esc


Purina®

IODINE CONCENTRATE 3.5

**WARNING: KEEP OUT OF REACH OF CHILDREN
CAUSES SKIN AND EYE IRRITATION
SEE OTHER WARNINGS ON SIDE PANEL**

Active Ingredients:
Alpha (p-nonylphenyl)-omega-hydroxypoly (oxyethylene)-
iodine complex (providing 3.5% titratable iodine) 27.60%
Phosphoric Acid 9.00%
Inert Ingredients: 63.40%
U.S. Patent Nos. 2,931,777-3,028,299

ACCEPTED
602-244
JAN 13 1973
UNDER FEDERAL INSECTICIDE
FUNGICIDE AND WEEDKILLER ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 602-244



TM

PU
ver
bac
Sal
Tri
tub
Sta
Sal
Esc

**Chemical
Specialties™**


NET CONTENTS: 1 GALLON

ma[®]

ATE 3.5

CH OF CHILDREN
RRITATION
SIDE PANEL

oxypoly (oxyethylene)-
atable iodine) 27.60%
..... 9.00%
..... 63.40%



TM

NET CONTENTS: 1 GALLON

ACCEPTED

602-244

JAN 15 1973

UNDE... INSECTICIDE
FUNGICIDE... FUNGICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. 602-244

PURINA IODINE CONCENTRATE 3.5 is a versatile germicide that kills many pathogenic bacteria and fungi such as *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Trichophyton interdigitale*, and *Mycobacterium tuberculosis*.

A.O.A.C. Use Dilution Confirmation Test

<i>Staphylococcus aureus</i>	1:512
<i>Salmonella choleraesuis</i>	1:512
<i>Pseudomonas aeruginosa</i>	1:512

A.O.A.C. In Vitro Test For Tuberculocidal Activity

<i>Mycobacterium tuberculosis</i>	1:256
-----------------------------------	-------

A.O.A.C. Fungicidal Test

<i>Trichophyton interdigitale</i>	Passes
-----------------------------------	--------

A.O.A.C. Germicidal Sanitizer Test
(1,000 ppm Hard Water)

<i>Staphylococcus aureus</i>	1:1536
<i>Escherichia coli</i>	1:1536

TAKE TIME



OBSERVE LABEL
DIRECTIONS

Manufactured by

RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63188