

17
2307459

SEP 13 1990

Ms. Jan Campbell
Purina Mill, Inc.
P.O. Box 66812
St. Louis, MS 63166-6812

Dear Ms. Jan Campbell:

Subject: Coumaphos Registration Standard - 2nd Round Review - Label
Revision
Purina Cattle Duster Insecticide
EPA Registration No. 602-250
Purina Lice and Fly Killer-CR
EPA Registration No. 602-251
Your Letter Dated January 31, 1990

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes indicated below prior to release for shipment of product bearing the amended labeling.

1. For both products make the following revisions:
 - a. Under **Note to Physician**, please insert the wording **"by injection"** in the phrase "Atropine sulfate by injection is antidotal."
 - b. Change the PESTICIDE DISPOSAL to the following:

Completely empty bag into application equipment. then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Submit five copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 15
Insecticide-Rodenticide Branch
Registration Division (H7505C)

(9-Square Logo[®])

Front Panel

PURINA[®]
BRAND

CATTLE DUSTER INSECTICIDE

FOR TRY OR BEEF CATTLE

EPA Reg. No. 602-250

3-Row Checkered[®]

FOR CONTROL OF HORNFLEES AND LICE
AIDS IN REDUCING FACE FLIES ON CATTLE

ACTIVE INGREDIENT:

0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) phosphorothioate	1%
INERT INGREDIENTS	99%
	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE OTHER PRECAUTIONS ON BACK PANEL

NET WT 12-1/2 LB

D-0873

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 13 1990

Under the Fungicide, and Herbicide Act, amended, for the pesticide registered under EPA Reg. No.

Manufactured By
PURINA MILLS, INC.
P. O. Box 66812

St. Louis, Missouri 63166-6812

NET 12-1/2 LB

PURINA[®]

BRAND

CATTLE DUSTER INSECTICIDE

Intended color of final label - yellow and white background with black type.
Type size of text - 3 Pt.

Back Panel

PURINA[®] CATTLE MUSTER INSECTICIDE
BRAND

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dusts. Avoid contact with eyes, skin, and clothing.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention if victim displays signs of poisoning.

If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS:

This product is toxic to birds, fish, and aquatic invertebrates. Do not contaminate water when disposing of equipment, washwater or sludge.

RESTRICTIONS:

For external, insecticidal use on above specified animals only. Avoid contamination of feed, troughs, water and water receptacles. Provide thoroughment of areas while dusting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

A ready-to-use insecticide dusting powder for use in the refillable PURINA Cattle Duster Bag -- a weather-proof bag for use in open or sheltered areas. This product can also be used for hand dusting. Effective in controlling hornflies and lice; aids in controlling face flies. No withdrawal period required; may be used with dairy cattle in production or beef cattle being fed for slaughter.

DUST BAG

Empty contents of bag (Cattle Duster Insecticide) into the Cattle Duster Bag through opening along edge of upper side. Close opening. Use 1 lb. of Cattle Duster Insecticide per 8 animals every two weeks. Do not exceed 3 lbs. of Cattle Duster Insecticide per animal per year.

HOW TO HANG:

Hang Cattle Duster Bag by means of rope or chain placed through straps along top edge. For Horn Fly and Lice control, adjust height so that bottom of bag will hang a little below the backline of the animals using it. To aid in reducing face flies, hang bag so that animals will be forced to dust face daily.

WHERE TO HANG:

Suspend Cattle Duster Bag in locations where animals will frequently dust themselves on a free-choice basis; or in gateways or lanes through which animals are forced to pass daily for water, feed, mineral.

Dairy Cattle: For lactating dairy cows, hang in the exit of the milking barn.

Restrictions: Do not hang Cattle Duster Bag over feed, mineral or water troughs.

HAND DUSTING

Apply not more than 2 oz. (6 level tablespoonfuls) evenly into the hair over the head, neck, shoulders, back and tailhead. Repeat as necessary, but not more often than every 14 days. (Maximum Seasonal Application Rate -- 26 applications/year)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

QUALITY CONTROLLED BY PURINA RESEARCH
EPA Est. No. 699-MO-1

8911 D

Printed in U.S.A.