

Purina LICE AND FLY KILLER-CR: A ready-to-use dusting powder for dairy or beef cattle and swine. Effective against lice and hornflies. No interval is required between treatment and use of meat or milk as food.

HOW TO USE

Animal	Insect	Directions
Cattle (Dairy or Beef)	Lice Hornflies	DIRECT APPLICATION: Dust evenly into the hair over the head, neck, shoulders, back and tail-head. Apply not more than 2 oz.* per animal. Repeat as necessary.
Swine	Lice	DIRECT APPLICATION: Dust evenly into the hair coat of shoulders and back. Apply not more than 1 oz.* per animal. Repeat as necessary but not more often than once every 10 days. BEDDING TREATMENT: Apply 2 oz.* uniformly over each 30 square feet of fresh, dry bedding. Repeat as necessary, but not more often than once every 10 days. DIRECT APPLICATION AND BEDDING TREATMENT: In severe infestations, both individual animals and the bedding may be treated as directed above.

*NOTE: 1 oz. equals approximately 3 level tablespoonfuls of LICE AND FLY KILLER-CR

RESTRICTIONS

For external, insecticidal use on above specified animals only. Avoid contamination of feed, troughs, water, and water utensils. Provide thorough ventilation while dusting.

CAUTIONS

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Exposure will result in cholinesterase depression.

To Physician—Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not reuse empty container for any other purpose. Destroy by burying or discard in safe place.

ENVIRONMENTAL HAZARD:


This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Use this product only as specified on this label.

QUALITY CONTROLLED BY PURINA RESEARCH
EPA Est. 602-MO-1



E.P.A. Reg. No. 602-251-AA



PURINA 



602-251

LICE & FLY KILLER-CR

FOR CONTROL OF LICE AND HORNFLIES ON DAIRY AND BEEF CATTLE AND LICE ON SWINE



Active Ingredient:
O-O-Diethyl O-(3-chloro-4-methyl-2-oxo-(2H)-1-benzopyran-7-yl) phosphorothioate 1%
Inert Ingredients: 99%
100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTIONS ON BACK PANEL

NET WT.: 16 OZ. (1 LB.)

Manufactured By
RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63188

Purina LICE AND FLY KILLER-CR: A ready-to-use dusting powder for dairy or beef cattle and swine. Effective against lice and hornflies. No interval is required between treatment and use of meat or milk as food.

HOW TO USE

Animal	Insect	Directions
Cattle (Dairy or Beef)	Lice Hornflies	DIRECT APPLICATION: Dust evenly into the hair over the head, neck, shoulders, back and tail-head. Apply not more than 2 oz.* per animal. Repeat as necessary.
Swine	Lice	DIRECT APPLICATION: Dust evenly into the hair coat of shoulders and back. Apply not more than 1 oz.* per animal. Repeat as necessary but not more often than once every 10 days. BEDDING TREATMENT: Apply 2 oz.* uniformly over each 30 square feet of fresh, dry bedding. Repeat as necessary, but not more often than once every 10 days. DIRECT APPLICATION AND BEDDING TREATMENT: In severe infestations, both individual animals and the bedding may be treated as directed above.

*NOTE 1 oz. equals approximately 3 level tablespoonfuls of LICE AND FLY KILLER-CR

RESTRICTIONS

For external, insecticidal use on above specified animals only. Avoid contamination of feed, troughs, water, and water utensils. Provide thorough ventilation while dusting.

CAUTIONS

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Exposure will result in cholinesterase depression.

To Physician—Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not reuse empty container for any other purpose. Destroy by burying or discard in safe place.


ENVIRONMENTAL HAZARD:


This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Use this product only as specified on this label

QUALITY CONTROLLED BY PURINA RESEARCH
EPA Est 602-MO-1





E.P.A. Reg. No. 602-251-AA

PURINA  TM



LICE & FLY KILLER-CR

**FOR CONTROL OF LICE AND HORNFLIES
ON DAIRY AND BEEF CATTLE
AND LICE ON SWINE**



Active Ingredient:
O-O-Diethyl O-(3-chloro-4-methyl-2-oxo-(2H)-
1-benzopyran-7-yl) phosphorothioate 1%
Inert Ingredients: 99%
100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTIONS ON BACK PANEL

NET WT.: 16 OZ. (1 LB.)

Manufactured By
RALSTON PURINA COMPANY
General Offices • Checkerboard Sq. • St. Louis, Mo. 63188