


E.P.A. Reg. No.
602-255-AA

NEW



Purina Rat Control Pellets



KILLS RATS and MICE

Active Ingredient:

Warfarin (3-(alpha-Acetylbenzyl)
4-hydroxycoumarin) 0.025%

Inert Ingredients 99.975%

100.000%

ACCEPTED
MAY 17 1975
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 602-255

CAUTION:

Keep Out of Reach of Children
See Other Cautions on Back Panel

NET WEIGHT 5 LBS.

Manufactured by
RALSTON PURINA COMPANY

General Offices, Checkerboard Square, St. Louis, Missouri 63188

DIRECTIONS FOR USE

Place RAT CONTROL PELLETS in locations where rats and mice feed, water, and/or travel. It is desirable to place the pellets along walls, in corners, and in concealed locations where rats and mice can feed unobserved. Contaminated or spoiled pellets should be replaced promptly. Where a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and the bait replenished as needed.

FOR RATS (Roof and Norway):

Use 4 to 16 ounces of RAT CONTROL PELLETS per placement. An uninterrupted supply of bait should be maintained for at least 10 days.

FOR HOUSE MICE:

Use tablespoon amounts ($\frac{1}{4}$ to $\frac{1}{2}$ ozs.) of RAT CONTROL PELLETS placed at 8 to 12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

POULTRY:

In caged houses, place bait along edge of dropping pits, walls, and in corners. For floor operations, bait buildings when depopulated and cleaned prior to putting down birds.

BARN, FEEDLOTS, KENNELS, RABBITRY, FUR FARMS, GAME BIRD OPERATIONS:

Place bait in protected areas outside of pens where animals or birds cannot reach it. Place bait in drain tile, under box, or under a board placed at an angle to building to protect from weather.

LABORATORY ANIMALS:

Place bait in corner of caged animal rooms, feed rooms, and equipment storage areas.

HOUSEHOLD:

Place bait out of reach of children and pets in protected areas of basements, garages, under porches and slabs, sewers, privies, and storage areas.

ENVIRONMENTAL HAZARDS:

Baits should be placed in locations not accessible to children, pets, wildlife, and domestic animals or in tamper-proof bait boxes. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CAUTION:

RAT CONTROL PELLETS contain an anticoagulant chemical which, if taken continuously by humans, domestic animals, or pets may reduce the clotting ability of the blood. Internal hemorrhage may result. Place bait out of reach of children, pets, and livestock. Should poisoning occur, give a tablespoonful of salt in a glass of warm water and repeat until vomit is clear. Call a physician immediately.

NOTE FOR PHYSICIAN:

When a human being has eaten this product, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of blood anticoagulants.

Do not reuse container. Destroy when empty.

QUALITY CONTROLLED BY PURINA RESEARCH