


PM16 602-255



**Purina**  
BRAND

# RAT CONTROL PELLETS



**KILLS RATS and MICE**

ACTIVE INGREDIENT:	
Warfarin, 3-(alpha-Acetylbenzyl)-	
4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
	100.000%

**ACCEPTED**

APR 10 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 602-255

KEEP OUT OF REACH OF CHILDREN  
**CAUTION:**  
 SEE BELOW FOR PRECAUTIONARY STATEMENTS

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Keep away from humans, domestic animals and pets. If swallowed this material may reduce the clotting ability of the blood and cause bleeding.

**NOTE FOR PHYSICIAN:** If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

**ENVIRONMENTAL HAZARDS:** Keep out of lakes, streams or ponds.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:** For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Place **RAT CONTROL PELLETS** in locations where rats and mice feed, water, and/or travel. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rats and mice or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

**FOR RATS (Roof and Norway):**

Use 4 to 16 ounces of **RAT CONTROL PELLETS** per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**FOR HOUSE MICE:**

Use ¼ to ½ ounce (1-2 level tablespoons) of **RAT CONTROL PELLETS** at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**POULTRY:**

In caged houses, place bait along edge of dropping pits, walls, and in corners. For floor operations, bait buildings when depopulated and cleaned prior to putting down birds.

832 H

**BARNs, FEEDLOTS, KENNELS, RABBITRY, FUR FARMS, GAME BIRD OPERATIONS:**

Place bait in protected areas outside of pens where animals or birds cannot reach it. Place bait in drain tile, under box, or under a board placed at an angle to building to protect from weather.

**LABORATORY ANIMALS:**

Place bait in corner of caged animal rooms, feed rooms, and equipment storage areas.

**HOUSEHOLD:**

Place bait out of reach of children and pets in protected areas of basements, garages, under porches and slabs, sewers, privies, and storage areas.

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent re-infestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present (nearby dumps), establish permanent bait stations and replenish bait as needed.

**DISPOSAL**  
 DO NOT REUSE EMPTY CONTAINER.  
 WRAP CONTAINER AND  
 PUT IN TRASH COLLECTION.

QUALITY CONTROLLED BY PURINA RESEARCH  
 EPA Reg. No. 602-MO-1  
 D-8307

Manufactured by  
**RALSTON PURINA COMPANY**  
 General Offices • One Centerville Square  
 St. Louis, Missouri 63103



EPA Reg. No. 602-255

Printed in U.S.A.

# NET WT 50 LB

10/21