

DANGER. CONTENTS UNDER PRESSURE
KEEP OUT OF REACH OF CHILDREN
READ CAREFULLY OTHER
CAUTIONS ON THE BACK PANEL

NET WT. 11 OZ.

Distributed by
RALSTON PURINA COMPANY

HOW TO USE PURINA FACEFLY BOMB

FACE FLIES: Remove any mucous secretions or saliva deposits from face of dairy cattle, beef cattle, and horses. Spray lightly, face and other body regions where face flies gather—4 to 5 seconds per head. Pay particular attention to the areas around the eyes and nostrils. Treat dairy cattle daily, preferably while they are stanchioned for morning milking.

For temporary protection of dairy cattle, beef cattle, and horses from horse flies, horn flies, stable flies, deer flies, house flies, mosquitoes, and gnats. Hold container 12 inches from animal. Apply lightly to hair coat with a sweeping motion. Apply to back, belly, ear, or leg regions where the specific insect attacking the animal usually feeds. Do not spray entire body area. Spray young animals sparingly. Do not apply to back of horses before they are saddled. Repeat as necessary for continued protection. Avoid wetting hide.

CAUTION: Harmful or fatal if swallowed. If swallowed, do not induce vomiting. Call physician immediately. Avoid contamination of feed and foodstuffs.

Do not contaminate milk or milk handling equipment. Do not spray or store near heat or open flame.

DANGER! Extremely flammable, contents under pressure. Do not use near fire, heated surfaces, sparks, or flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

753B D-3634 Printed in U.S.A.

EPA Est. 11715-TN-1

ACCEPTED

MAY 15 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 602-256