

## HOW TO USE PURINA FLY SUGAR BAIT

Scatter Purina Fly Sugar Bait daily or as needed to reduce fly population. Distribute bait from container or with jar or can to avoid handling bait. Keep Fly Sugar Bait in original container tightly closed and store in cool place.

**Loading Sheds, Livestock Pens, Pig Parlors, Stables, Broiler Houses:** Use only around outside areas. Scatter Fly Sugar Bait at the rate of ¼ lb. per 1000 sq. ft. on ground areas where flies congregate. Ideal baiting sites are around doorways, along fence lines, soiled areas, or in alleyways of feedlots.

**Caged Poultry and Turkeys on Wire:** Scatter Fly Sugar Bait on walkways at the rate of ¼ lb. per 500 sq. ft. to aid in controlling house flies. Also scatter outdoors around doorways.

**Dog Kennels and Furbearing Animal Units:** Scatter Fly Sugar Bait around outside areas of dog pens, runways, and other surfaces out of reach of the animals. For cage units, scatter on walkways at the rate of ¼ lb. per 500 sq. ft.

**Food, Poultry and Meat Processing Plants:** Scatter Fly Sugar Bait at the rate of ¼ lb. per 1000 sq. ft. in inedible departments only. Fly Sugar Bait may be used outside livestock pens or loading and unloading docks.

### WARNING

May be harmful or fatal if swallowed,  
inhaled or absorbed through skin.

**DO NOT** get on skin, in eyes, or on clothing. Wash thoroughly with soap and water after using. Keep out of reach of children.

Do not contaminate feed and foodstuffs. Do not apply where poultry or animals, especially young calves, can pick it up or lick it. Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Use only in areas **inaccessible** to food producing animals.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Use this product only as specified on this label.

Do not re-use empty container. Destroy container by perforating and bury or discard in a safe place.

Should illness occur, get prompt medical aid.

**NOTE TO PHYSICIAN:** Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

**ANTIDOTE:** If swallowed induce vomiting with 1 tablespoon of salt and warm water.

757 B

EPA Est. 602-MO-1 Printed in U.S.A.

# D-3753

QUALITY CONTROLLED BY PURINA RESEARCH

E.P.A. Reg. No. 602-257-AA



# PURINA®

## FLY SUGAR BAIT

Attracts and Kills House Flies



### Active Ingredient:

\*Dimethyl 3-hydroxyglutaconate dimethyl phosphate + . . . . . 1.0%

Inert Ingredients . . . . . 99.0%

100.0%

\*Bomyl® Reg. TM Allied Chemical Corporation  
+U. S. Patent No. 2,891,887

### WARNING: KEEP OUT OF REACH OF CHILDREN

See Other Warnings on Back Panel

KEEP CONTAINER CLOSED  
STORE IN A COOL DRY PLACE

### NET WT.: 16 OZ. (1 lb.)

Manufactured by  
RALSTON PURINA COMPANY  
General Offices  
Checkerboard Square  
St. Louis, Missouri 63188