

## HOW TO USE PURINA FLY BAIT GRITS

Scatter Purina Fly Bait Grits daily or as needed to reduce fly population. Distribute bait from container or with jar or can to avoid handling bait. Keep Fly Bait Grits in original container tightly closed and store in cool place.

**Loafing Sheds, Livestock Pens, Pig Parlors, Stables, Broiler Houses:** Use only around outside areas. Scatter Fly Bait Grits at the rate of 1/4 lb. per 1000 sq. ft. on ground areas where flies congregate. Ideal baiting sites are around doorways, along fence lines, soiled areas, or in alleyways or feedlots.

**Caged Poultry and Turkeys on Wire:** Scatter Fly Bait Grits on walkways at the rate of 1/4 lb. per 500 sq. ft. to aid in controlling house flies. Also scatter outdoors around doorways.

**Dog Kennels and Furbearing Animal Units:** Scatter Fly Bait Grits around outside areas of dog pens, runways, and other surfaces out of reach of the animals. For cage units, scatter on walkways at the rate of 1/4 lb. per 500 sq. ft.

**Food, Poultry and Meat Processing Plants:** Scatter Fly Bait Grits at the rate of 1/4 lb. per 1000 sq. ft. in inedible departments only. Fly Bait Grits may be used outside livestock pens or loading and unloading docks.

### WARNING

May be harmful or fatal if swallowed, inhaled or absorbed through skin.  
DO NOT get on skin, in eyes, or on clothing. Wash

thoroughly with soap and water after using. Keep out of reach of children.

Do not contaminate feed and fordstuffs. Do not apply where poultry or animals, especially young calves, can pick it up or lick it. Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Use only in areas **inaccessible** to food producing animals.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Use this product only as specified on this label.


Do not re-use empty container. Destroy container by perforating and bury or discard in a safe place.

Should illness occur, get prompt medical aid.

**NOTE TO PHYSICIAN:** Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

**ANTIDOTE:** If swallowed induce vomiting with 1 tablespoon of salt and warm water

E. P. A. Reg. No.  
602-258-AA



TM

**PURINA®**

# FLY BAIT GRITS

ATTRACTS and KILLS HOUSE FLIES



### Active Ingredient:

\*Dimethyl 3-hydroxyglutaconate dimethyl phosphate+ . . . . . 1.0%

Inert Ingredients . . . . . 99.0%

100.0%


\*Bomyl® Reg. TM Allied Chemical Corporation  
+ U. S. Patent No. 2,891,887

### WARNING

KEEP OUT OF REACH OF CHILDREN  
See Other Warnings on Back Panel  
KEEP CONTAINER CLOSED  
STORE IN A COOL DRY PLACE

**NET WT. 2 lbs.**

Manufactured by  
**RALSTON PURINA COMPANY**  
General Offices  
Checkerboard Square  
St. Louis, Missouri 63188



**D-3733**

(EPA Est. 602 MO-1)

Printed in U. S. A.

**QUALITY CONTROLLED BY PURINA RESEARCH**