

602-260

08/16/1977

**PRECAUTIONARY STATEMENTS
CAUTION**

HAZARD TO HUMANS AND DOMESTIC ANIMALS: Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Atropine Sulfate is antidotal. If swallowed induce vomiting with 1 tablespoonful of salt in a glass of warm water.

Do not contaminate feed and food-stuffs. Do not apply where poultry or animals, especially young calves, can pick it up or lick it. Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Use only in areas inaccessible to food producing animals.

Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to wildlife and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Lot Nbr.: _____ Exp. Date: _____
7610 C

Scatter Purina Fly Patrol (Fly Bait) daily or as needed to reduce fly population. Distribute bait from container or with jar or can to avoid handling bait. Keep Fly Patrol in original container tightly closed and store in cool place.

Loafing Sheds, Livestock Pens, Pig Parlors, Stables, Broiler Houses: Use only around outside areas. Scatter Fly Patrol at the rate of 1/4 lb. per 1000 sq. ft. on ground areas where flies congregate. Ideal baiting sites are around doorways, outside of fence lines, soiled areas, or in alleyways of feedlots. Place bait on clean paper if floors are lined.

Caged Poultry and Turkeys on Wire: Scatter Fly Patrol on walkways at the rate of 1/4 lb. per 500 sq. ft. to aid in controlling house flies. Also scatter on floors around doorways.

Dog Kennels and Furbearing Animal Units: Scatter Fly Patrol around outside areas of dog pens, runways, and other surfaces, out of reach of the animals. For cage units, scatter on walkways at the rate of 1/4 lb. per 500 sq. ft.

Food, Poultry and Meat Processing Plants: Scatter Fly Patrol at the rate of 1/4 lb. per 1000 sq. ft. in receiving departments only. Fly Patrol may be used outside livestock pens or loafing sheds.

STORAGE AND DISPOSAL:

Do not store near food or feed.
Do not re-use empty container.

Destroy container and bury or discard in a safe place away from streams, drainage or water supplies.

E.P.A. Est. No. 602-MO-1


Printed in U.S.A.

QUALITY CONTROLLED BY PURINA RESEARCH



**PURINA[®]
FLY PATROL[™]
(FLY BAIT)**

Attracts and Kills House Flies



Active Ingredient:	Methomyl (s-methyl N-[methylcarbamoyl] oxy) thioacetimidate	1.0%
Inert Ingredients	99.0%
		100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR PRECAUTIONARY STATEMENTS**

NET WT. 16 OZ. (1 LB.)

Manufactured By
RALSTON PURINA COMPANY
GENERAL OFFICES • CHECKERBOARD SQUARE
St. Louis, Missouri 63188

ACCEPTED
AUG 16 1977
EPA REG. NO. 602-260
Federal Insecticide, Rodenticide Act

602-260

11