

Reason For Issue: Updated
per EPA Reregistration Standard

Date Issued: 6/20/89 (Yellow)

Prev. Date Issued: 2/17/86

(9-Square Logo[®])

Front Panel

(3-Row Checkerband[®])

PURINA[®]

EPA Reg. No. 602-260

FLY PATROL[®]
BRAND

(Fly Bait)

Attracts and Kills House Flies

(10 Pt. Type Size)

INGREDIENT STATEMENT

ACTIVE INGREDIENT:

Methomyl (S-Methyl-N-[(methylcarbamoyl)oxy]-thioacetimidate . . .	1.0%
INERT INGREDIENTS	99.0%
	100.0%

KEEP OUT OF REACH OF CHILDREN

--(10 Pt. Type Size)


CAUTION

--(14 Pt. Type Size)

SEE BACK PANEL FOR PRECAUTIONARY STATEMENTS

--(10 Pt. Type Size)

NET WT 16 OZ (1 LB)




D-3701

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812

St. Louis, Missouri 63166-6812 U.S.A.

NET 16 OZ

PURINA[®] FLY PATROL[®]
BRAND



(Intended color of final label)

Front Panel-Yellow & White background with
black type

Back Panel-White background with black type
Type Size for Directions For Use - 6 Pt.

Back Panel

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes or skin with plenty of water.

Get medical attention if irritation persists. Antropine Sulfate is antidotal.

PRACTICAL TREATMENT: If swallowed: Call a physician or Poison Control Center.

Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

Do not contaminate feed and foodstuffs. Do not apply where poultry or other animals, especially dogs and young calves, can pick it up or lick it.

Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Use only in areas inaccessible to food producing animals.

Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas when food is present.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to birds and fish. Collect, cover or incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PERSONAL PROTECTIVE CLOTHING

Use only when wearing the following personal protective clothing during loading, application, repairing and cleaning of application equipment, and disposal of the pesticide: long-sleeve shirt; long-legged pants; shoes and socks; gloves.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water.

ALWAYS WASH HANDS, FACE, AND ARMS WITH SOAP AND WATER BEFORE USING TOBACCO PRODUCTS, EATING, DRINKING, OR TOILETING.

Scatter PURINA FLY PATROL (Fly Bait) daily or as needed to reduce fly population. Distribute bait from container or with jar or can to avoid handling bait. Keep FLY PATROL in original container tightly closed and store in cool place.

Loafing Sheds, Livestock Pens, Pig Parlors, Stables, Broiler Houses. Use only around outside areas. Scatter FLY PATROL at the rate of $\frac{1}{2}$ lb per 1000 sq. ft. on ground areas where flies congregate. Ideal baiting sites are around doorways, outside of fence lines, soiled areas, or in alleyways of feedlots. Place bait on clean paper if floors are limed.

Caged Poultry and Turkeys on Wire: Scatter FLY PATROL on walkways at the rate of $\frac{1}{4}$ lb per 500 sq. ft. to aid in controlling house flies. Also scatter outdoors around doorways.

Dog Kennels and Furbearing Animal Units: Scatter FLY PATROL around outside areas of dog pens, runways, and other surfaces out of reach of the animals. For cage units, scatter on walkways at the rate of $\frac{1}{4}$ lb. per 500 sq. ft.

Food, Poultry and Meat Processing Plants: Scatter FLY PATROL at the rate of $\frac{1}{4}$ lb per 1000 sq. ft. in inedible department only. FLY PATROL may be used outside livestock pens or loading and unloading docks.

STORAGE AND DISPOSAL

--(10 Pt. Type Size)

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area in original container.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay of smoke.

QUALITY CONTROLLED BY PURINA RESEARCH

EPA Est. No. 602-MO-1
44946-MO-1 T

Lot No. preceded by T indicates
EPA Est. No. 44946-MO-1

8906 J

Printed in U.S.A.

