

PM16 602-266

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling
Use restrictions: For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings and similar man-made structures.
Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.
Selection of treatment areas: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls by gnawed openings in or beside burrows, in corners and concealed places, between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.
Application directions: RATS: Apply a minimum of one PLACE PACK RAT Control per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days. MICE: Apply



Purina

PLACE PACK RAT CONTROL



Kills Norway Rats, Roof Rats and House Mice

ACTIVE INGREDIENT
Warfarin (3-(alpha-Acetylbiphenyl)-4-hydroxycoumarin)
INERT INGREDIENTS

REREGISTERED

ACCEPTED

DEC 20 1982

Under Federal FIFRA
Fungicide, Insecticide and Rodenticide Act
as amended, for the pest
registered under
EPA Reg. No. 602-266

0.025%
94.975%
100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BELOW FOR PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: Keep out of lakes, streams or ponds.

NET WT 4 OZ

RALSTON PURINA COMPANY
Rat Control Pack
St. Louis, Missouri 63104



Printed in U.S.A.

D-8324

628 F

602-266

EPA Reg. No.

BY PURINA RESEARCH
QUALITY CONTROLLED

STORAGE AND DISPOSAL: Do not re-use empty container. Wrap container and put in trash collection.
Replace all contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.
one PLACE PACK RAT Control at 8 to 12 foot intervals. More placements may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

1/01