


E.P.A. REG. NO. 602-267



PURINA® DAIRY MINERAL FLY LARVICIDE (Mineral Feed)



For Free-Choice Feeding to Dairy Cattle

To Prevent the Development of Horn Flies, Face Flies, House Flies and Stable Flies in Manure of Treated Cattle.

| | |
|--|-----------|
| Active ingredient: | By Weight |
| 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate..... | 1.45% |
| Inert ingredients..... | 98.55% |
| | 100.00% |

GUARANTEED FEED ANALYSIS

| | |
|-----------------------------------|--------|
| Calcium (Ca) not less than..... | 16.0 % |
| Calcium (Ca) not more than..... | 19.0 % |
| Phosphorus (P) not less than..... | 16.0 % |
| Iodine (I) not less than..... | 0.016% |

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating and smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE GENERAL USE CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Feed Purina Dairy Mineral Fly Larvicide from a sheltered feeder as the only free-choice source of mineral. Cattle should consume an average of 0.011 lbs. of Purina Dairy Mineral Fly Larvicide per 100 lbs. of body weight per day (1 1/4 oz./1000 lbs. body wt./day). This will supply the recommended average daily intake of 70 mg RABON insecticide per 100 lbs. body weight. Animals should not be starved for minerals before using this product. Place feeder near watering or loafing areas. Reduce or increase the number of feeding locations or relocate if consumption is above or below recommended daily intake. Provide plain salt free-choice since Dairy Mineral Fly Larvicide contains no salt.

Start feeding Purina Dairy Mineral Fly Larvicide early in the spring before flies appear and continue feeding until cold weather restricts fly activity. Resume feeding Purina Dairy Mineral until spring.

Purina Dairy Mineral Fly Larvicide may be fed up to slaughter and to lactating dairy cows. Proper consumption prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplement fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, use this product in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not store near food or feed. Do not re-use empty container. Destroy container and bury or discard in a safe place away from streams, drainage or water supplies.

ACCEPTED

JUN 14 1979

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL
FOR ADDITIONAL PRECAUTIONS**

RABON® INSECTICIDE Reg. T.M. Shell Chemical Company

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 602-267

Manufactured By
RALSTON PURINA COMPANY
GENERAL OFFICES - CHECKERBOARD SQUARE
St. Louis, Missouri 63188

QUALITY-CONTROLLED BY PURINA RESEARCH

EPA 20.910. 609110-1

793 B

Printed in U.S.A.

NET WT. 50 LBS.

D-2218

602-267

6-14-1979