


602-271 10F2
pm 15

PURINA[®]

FRESHSTART D & H

FLY LARVICIDE BLOCK

(Mineral Block)

**PRECAUTIONARY STATEMENTS****CAUTION**

HAZARDS TO HUMANS: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating and smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Feed free choice in weather-protected feeders. The ratio of the number of blocks to animal units will help control consumption to the desired 2 pounds or 500 lbs. of body weight per day. This will supply the recommended average intake of 70 mg RABON per hundred pounds of body weight. Animals should not be starved for minerals before using this product. Do not feed free choice sources of salt or mineral containing salt in any form. Place blocks near watering and loafing areas.

Start feeding FRESHSTART D & H FLY LARVICIDE BLOCKS early in the spring before flies appear and continue feeding until cold weather restricts fly activity. Resume feeding FRESHSTART D & H BLOCK until spring.

Purina Freshstart D & H Fly Larvicide Blocks may be fed up to slaughter and to lactating dairy cows.

Proper consumption prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, use this product in conjunction with other good management and sanitation practices.

To Prevent the Development of Horn Flies, Face Flies, House Flies and Stable Flies in Manure of Treated Cattle.

Active Ingredient:	By Weight
*2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.055%
Inert Ingredients	99.945%
	100.00%

FEED INGREDIENTS

Plant protein products, processed grain by-products, forage products, grain products, cane molasses, urea, defluorinated phosphate, calcium carbonate, sulfur, iodized salt, vitamin A supplement, ethoxy quin (a preservative), D activated animal sterol, magnesium oxide, potassium chloride, iron oxide, manganous oxide, copper oxide, cobalt carbonate, zinc oxide.

GUARANTEED ANALYSIS

Crude Protein not less than	28.0 %
Include not more than 13.5% equivalent protein from non-protein nitro. ..	
Crude fat not less than	0.5 %
Crude fiber not more than	10.0 %
Calcium (Ca) not less than	2.5 %
Calcium (Ca) not more than	3.5 %
Phosphorus (P) not less than	1.5 %
Iodine (I) not less than	0.0005%
Magnesium (Mg) not less than	1.0 %
Salt (NaCl) not less than	12.5 %
Salt (NaCl) not more than	15.0 %
Vitamin A not less than	30,000 IU/lb
Vitamin D not less than	10,000 IU/lb

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

KEEP OUT OF REACH OF CHILDREN**CAUTION.**

See Below For Additional Precautionary Statements

*RABON[®] INSECTICIDE Reg. T.M. Shell Chemical Company

Manufactured By
RALSTON PURINA COMPANY
GENERAL OFFICES - CHECKERBOARD SQUARE

QUALITY CONTROLLED BY PURINA RESEARCH
EPA Est. No. 602-MO-1

ACCEPTED

JUN 14 1979

CAUTION

HAZARDS TO HUMANS: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating and smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Feed free choice in weather-protected feeders. The ratio of the number of blocks to animal units will help control consumption to the desired 2 pounds per 500 lbs. of body weight per day. This will supply the recommended average intake of 70 mg RABON per hundred pounds of body weight. Animals should not be starved for minerals before using this product. Do not feed free choice sources of salt or mineral containing salt in any form. Place blocks near watering and loafing areas.

Start feeding FRESHSTART D & H FLY LARVICIDE BLOCKS early in the spring before flies appear and continue feeding until cold weather restricts fly activity. Resume feeding FRESHSTART D & H BLOCK until spring.

Purina Freshstart D & H Fly Larvicide Blocks may be fed up to slaughter and to lactating dairy cows.

Proper consumption prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, use this product in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

QUALITY CONTROLLED BY PURINA RESEARCH
EPA Est. No. 602-MO-1


795 C

ACCEPTED

JUN 14 1979

Under Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 602-271

FRESHSTART D & H FLY LARVICIDE BLOCK (Mineral Block)



To Prevent the Development of Horn Flies, Face Flies, House Flies and Stable Flies in Manure of Treated Cattle.

Active Ingredient:	By Weight
*2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphorothioate	0.055%
Inert Ingredients	99.945%
	100.00%

FEED INGREDIENTS

Plant protein products, processed grain by-products, forage products, grain products, cane molasses, urea, defluorinated phosphate, calcium carbonate, sulfur, iodized salt, vitamin A supplement, ethoxyquin (a preservative), D activated animal sterol, magnesium oxide, potassium chloride, iron oxide, manganous oxide, copper oxide, cobalt carbonate, zinc oxide.

GUARANTEED ANALYSIS

Crude Protein not less than	28.0	%
Include not more than 13.5% equivalent protein from non-protein nitrogen		
Crude fat not less than	0.5	%
Crude fiber not more than	10.0	%
Calcium (Ca) not less than	2.5	%
Calcium (Ca) not more than	3.5	%
Phosphorus (P) not less than	1.5	%
Iodine (I) not less than	0.0005%	
Magnesium (Mg) not less than	1.0	%
Salt (NaCl) not less than	12.5	%
Salt (NaCl) not more than	15.0	%
Vitamin A not less than	30,000 IU/lb	
Vitamin D not less than	10,000 IU/lb	

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Below For Additional Precautionary Statements

*RABON® INSECTICIDE Reg. T.M. Shell Chemical Company

Manufactured By
RALSTON PURINA COMPANY
GENERAL OFFICES - CHECKERBOARD SQUARE
St. Louis, Missouri 63108

NET WT. 33 1/3 LBS.

D-383