

PURINA SUP-R-LIX 16 .303 PTZ

Liquid Feed Supplement with Phenothiazine

For Ruminants Only

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle.

Licensed Under U. S. Pat. 3901976

ACTIVE INGREDIENT:	
Phenothiazine	0.303%
INERT INGREDIENTS:	99.697%
	100.000%

Guaranteed Analysis

Crude protein not less than (This includes not more than 14.0% equivalent protein from non-protein nitrogen)	16.0%
Crude fat not less than	5.0
Calcium (Ca) not less than	0.5
Calcium (Ca) not more than	1.5
Phosphorus (P) not less than	0.8
Iodine (I) not less than	0.00005
Vitamin A not less than	7,500 IU/lb.
Vitamin D not less than	1,500 IU/lb.

Feed Ingredients: Cane molasses, urea, ammonium polyphosphate solution, animal fat preserved with BHA, calcium carbonate, attapulgite clay, vitamin A supplement, D activated animal sterol, manganese sulfate, iron sulfate, copper sulfate, ethylene diamine dihydriodide, cobalt sulfate, zinc sulfate.

PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

CAUTION

HAZARDS TO HUMANS: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating and smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL: Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Feed at a rate of 1 lb. to animals weighing 550 lbs. body weight. Each animal should receive .25 g phenothiazine per 100 lbs. body weight (1.375 g/hd/day).
2. Provide a source of all the good forage the cattle will consume. During periods of extreme shortage of forage, it is recommended that hay or silage also be fed.
3. Provide an unlimited supply of clean, fresh water near the feeding areas.
4. Feed amounts only as directed per head per day. Observe cattle closely for the first 8 days to be sure that daily intake requirement is not exceeded and that Sup-R-Lix is not overconsumed. If consumption of Sup-R-Lix 16 is higher than desired, switch to Purina Sup-R-Lix Breeder 24 PTZ Liquid Feed Supplement.
5. Provide a source of mineral and salt free-choice.
6. Cattle must have consumed other dry feed prior to being fed Sup-R-Lix 16 PTZ for the first time.
7. Do not provide starved cattle free access to Sup-R-Lix 16 PTZ.
8. Do not feed Sup-R-Lix 16 in conjunction with other urea (NPH) containing feed, liquid or dry.
9. Non-approved or illegal additives are not to be added.
10. If feeder tank runs dry and is refilled, observe cattle closely for three days to prevent overconsumption.
11. Feed tanks should be kept clean and free from contamination of all kinds before refilling.

SUP-R-LIX 16 SHOULD ONLY BE FED ACCORDING TO THE ABOVE DIRECTIONS.

CAUTION: Do not feed to lactating dairy animals. Occasional sensitivity to Phenothiazine has been reported.

NOTE: This product kills horn fly and face fly maggots in manure but does not control existing adult flies. If necessary, spray cattle for maximum protection.

Because face flies migrate from farm to farm, it is essential that a larvicide program be employed on an area basis to assist in reducing the adult face fly population.

NET WEIGHT: BULK

EPA Est. No. 602-10-1

Manufactured By:
RALSTON PURINA COMPANY
General Offices
Checkerboard Square
St. Louis, Missouri 63108

Printed in U.S.A.

BEST DOCUMENT AVAILABLE

13