



PURINA

E.P.A. Reg. No. 602-289

# METHOXYCHLOR SPRAY

## EMULSIFIABLE INSECTICIDE CONCENTRATE



ACTIVE INGREDIENT **D-6050**

\*Methoxychlor Technical  
INERT INGREDIENTS

1,4-dimethoxychloro-2,2-bis(4-chlorophenyl)-1,1,1-trifluoroethane, 1,1,1-trifluoro-2,2-bis(4-methoxyphenyl)ethane, 1,1,1-trifluoro-2,2-bis(4-methoxyphenyl)ethane isomers and reaction products

**QUALITY CONTROLLED**  
AUG 04 1980  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, recommended for the pesticide registered under Reg. No. 602-289



**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
SEE BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS & DOMESTIC ANIMALS  
**CAUTION**  
Avoid contact with skin. Avoid breathing of spray mist. In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. Get medical attention if irritation persists.  
**ENVIRONMENTAL HAZARD**  
This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, streams, ponds. Do not apply when weather conditions favor runoff or drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**  
It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
**LIVESTOCK**—(except dairy animals)—For control of certain external parasites. *Use as spray only. Do not use in dipping vats.*  
**CATTLE**—Use 1 quart Purina Methoxychlor Spray to 12 gallons of water. Apply 2 quarts of spray per mature animal for control of hornflies. For control of short-horned and long-horned cattle lice, apply as a drenching spray.  
**HOGS**—For control of lice, use 1 quart Purina Methoxychlor Spray to 12 gallons of water applied as a drenching spray.  
**SELF APPLICATOR FOR FLY CONTROL ON CATTLE**—(except dairy animals)—For control of hornflies, use 1 gallon of Purina Methoxychlor Spray to 12 gallons of water.

Do not use in conjunction with kerosene, fuel or furnace type oil. Do not use any other type of oil. 1 gallon of solution to saturate burlap. Repeat treatments every two to three weeks.

**CONTACT OR SPACE SPRAY FOR FLIES:** To reduce populations of house flies and stable flies in farm buildings, in barnyards, in alleys, in areas adjacent to farm buildings, etc. Mix 1 pint of Purina Methoxychlor Spray with 2 gallons of water. Apply in space by mist blower or directly to flies by hydraulic sprayer, hand sprayer or cylindrical pressure sprayer. Avoid contamination of milk handling equipment, feed stuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying.

**GRAIN STORAGE BINS:** To clean up infestations in empty storage bins, use 1 gallon of Purina Methoxychlor Spray in 10 gallons of water. Apply at the rate of 2 gallons of spray per 1000 sq. ft. before products are stored. This aids in control of cadelle, flat grain beetle, saw-toothed grain beetle, lesser grain borer, red flour beetle, long-necked flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus beetle. **Note:** Do not add grain to bin for at least 24 hours or until walls have dried thoroughly.

**STORAGE AND DISPOSAL:**  
Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.  
**Pesticide Disposal:** Pesticide spray mixture or residue that cannot be used or chemically reprocessed **must** be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.  
**Container Disposal:** Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill or bury in a safe place.  
**General:** Consult Federal, State or Local Disposal Authorities for appropriate reuse procedures.

E.P.A. Est. No. 602-MO-1  
**QUALITY CONTROLLED BY PURINA RESEARCH**

**RALSTON PURINA COMPANY**  
R.D. 2, Box 100, St. Louis, Mo. 64503

**NET 5 GAL.**