

E.P.A. REG. NO. 602-



PURINA®

GENERAL SEVIN*-7

INSECTICIDE

Controls Garden and Ornamental Insects

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

NOTE FOR PHYSICIAN

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

CARROTS

LETTUCE

OKRA

TOMATO EGGPLANT PEPPER

POTATO

INTERVAL BEFORE HARVEST

CAUTION

ORNAMENTALS

LAWNS

EARWIGS

DISPOSAL

DIRECTIONS FOR USE

VEGETABLES

APPLICATION Apply with a spray nozzle to the foliage of plants. Do not apply to plants that are stressed or in bloom. Do not apply to plants that are to be eaten raw.

BEANS Apply to the foliage of beans. Do not apply to beans that are to be eaten raw.

CABBAGE COLLARDS BROCCOLI BRUSSELS SPROUTS CAULIFLOWER MUSTARD GREENS TURNIPS KOHLRABI AND SPINACH Apply to the foliage of these vegetables.

CUCUMBERS SQUASH CANTALOUPE PUMPKINS MELONS Apply to the foliage of these vegetables.

CORN Apply to the foliage of corn.

Apply to the foliage of plants. Do not apply to plants that are stressed or in bloom. Do not apply to plants that are to be eaten raw.

DUAL CONTROLLED BY PURINA RESEARCH

ACTIVE INGREDIENT
95.0% Deltamethrin (0.05% deltamethrin)
NET WEIGHT

9
100

*Trademark of the manufacturer of the active ingredient, deltamethrin.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE BACK PANEL

NET WT. 1 LB.

RALSTON PURINA COMPANY

1000 North Lincoln Street, St. Louis, Missouri 63103

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

NOTE FOR PHYSICIAN

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

DIRECTIONS FOR USE

VEGETABLES

CAUTION: Do not eat any vegetables treated with this insecticide until the specified interval has elapsed. Do not eat any vegetables treated with this insecticide until the specified interval has elapsed. Do not eat any vegetables treated with this insecticide until the specified interval has elapsed.

CABBAGE COLLARDS BROCCOLI BRUSSELS SPOUTS CAULIFLOWER MUSTARD GREENS TURNIPS KOLRABI AND SPINACH

PEPPERS SQUASH CANTALOUPE PUMPKINS

CARROTS

LETTUCE

ONIONS

TOMATOES EGGPLANT PEPPER

POTATOES

INTERVAL BEFORE HARVEST

CAUTION

ORNAMENTALS

LAWNS

EARWIGS

DISPOSAL




PURINA

GENERAL SEVIN*-7

INSECTICIDE

Controls Garden and Ornamental Insect Pests

E.P.A. REG. NO. 602-296



KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE ADDITIONAL PRECAUTIONARY STATEMENTS
ON THE BACK PANEL

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS
ON THE BACK PANEL

NET WT. 1 LB.

RALSTON PURINA COMPANY

QUALITY CONTROLLED BY PURINA RESEARCH

E.P.A. REG. NO. 602-21



PURINA[®]

GENERAL SEVIN*-7

INSECTICIDE

Controls Garden and Ornamental Insec

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

See back panel for full text of precautionary statements.

NOTE FOR PHYSICIAN

See back panel for full text of note for physician.

STATEMENT OF PRACTICAL TREATMENT

See back panel for full text of statement of practical treatment.

ENVIRONMENTAL HAZARDS

See back panel for full text of environmental hazards.

APPLES
LETTUCE
ORRA
TOMATO, EGGPLANT, PEPPER
POTATO
INTERVAL BEFORE HARVEST
CAUTION

ORNAMENTALS

LAWNS

EARWIGS

DISPOSAL

DIRECTIONS FOR USE

VEGETABLES

APPLICATION See back panel for full text of application instructions.

BEANS See back panel for full text of instructions for beans.


CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, MUSTARD GREENS, TURNIPS, KOHLRABI AND SPINACH See back panel for full text of instructions for these vegetables.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS See back panel for full text of instructions for these vegetables.

CORN See back panel for full text of instructions for corn.

See back panel for full text of additional directions for use.

QUALITY CONTROLLED BY PURINA RESEARCH



Active Ingredient: Cyfluthrin (0.05%)
Other Ingredients: 99.95%

Net Weight: 4 lb. (1.81 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE BACK PANEL

NET WT. 4 LB.

RALSTON PURINA COMPANY

GENERAL OFFICES • ONE KERR AVENUE
ST. LOUIS, MISSOURI 63164

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Irritation to eyes, nose, throat, and skin. Avoid contact with eyes, nose, throat, and skin. Wash thoroughly with soap and water after handling. Avoid breathing dust or mist. Avoid contact with clothing.

NOTE FOR PHYSICIAN: See back panel for first aid information.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, flush with water for at least 15 minutes. In case of skin contact, wash with soap and water. In case of inhalation, move to fresh air.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic life. Do not apply near water bodies. Do not apply to lawns or ornamentals.

DIRECTIONS FOR USE

Apply to foliage of plants and lawns. Do not apply to lawns or ornamentals.

VEGETABLES

APPLICATION: Begin applications when insects first observed and repeat weekly. Do not apply to lawns or ornamentals.

BEANS: To control Mexican Bean Beetle, apply at rate of 10 pounds per acre. To control Earwig, apply at rate of 10 pounds per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPOUTS, CAULIFLOWER, MUSTARD GREENS, TURNIPS, KOHLRABI AND SPINACH: To control Cabbage Root Fly, apply at rate of 10 pounds per acre.

CUMBERS, SQUASH, CANTALOUPE, PUMPKINS: To control Cucumber Beetle, apply at rate of 10 pounds per acre.

PEARS: To control Pear Leafhopper, apply at rate of 10 pounds per acre.

CARROTS

LETTUCE

OKRA

TOMATO, EGGPLANT, PEPPER

POTATO

INTERVAL BEFORE HARVEST

CAUTION

ORNAMENTALS

Do not apply to lawns or ornamentals. This product is highly toxic to fish and aquatic life.

LAWNS

Do not apply to lawns. This product is highly toxic to fish and aquatic life.

EARWIGS

To control Earwig, apply at rate of 10 pounds per acre. Do not apply to lawns or ornamentals.

DISPOSAL

Do not dispose of this product in water. Dispose of in a secure container.

QUALITY CONTROLLED BY PURINA RESEARCH




PURINA (R)

GENERAL SEVIN*-7

INSECTICIDE

Controls Garden and Ornamental Insect Pests

E.P.A. REG. NO. 602-296



ACTIVE INGREDIENT: Cyfluthrin, 1.5%
INERT INGREDIENTS: 98.5%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE BACK PANEL

NET WT. 4 LB.

RALSTON PURINA COMPANY

St. Louis, Missouri 63103

E.P.A. REG. NO. 602-296



PURINA[®]
BRAND

GENERAL SEVIN*-7

INSECTICIDE

Controls Garden and Ornamental Insect Pests



PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS—Irritant to eyes. Avoid contact with eyes. Wash thoroughly with water after use. Avoid contact with skin. Wash thoroughly with water after use. Avoid contact with clothing. Wash thoroughly with water after use. Avoid contact with food. Wash thoroughly with water after use. Avoid contact with children. Wash thoroughly with water after use. Avoid contact with pets. Wash thoroughly with water after use.

NOTE FOR PHYSICIAN

See back panel for first aid instructions.

STATEMENT OF PRACTICAL TREATMENT

Wash thoroughly with water after use. Avoid contact with eyes. Wash thoroughly with water after use. Avoid contact with skin. Wash thoroughly with water after use. Avoid contact with clothing. Wash thoroughly with water after use. Avoid contact with food. Wash thoroughly with water after use. Avoid contact with children. Wash thoroughly with water after use. Avoid contact with pets. Wash thoroughly with water after use.

ENVIRONMENTAL HAZARDS

See back panel for environmental hazard information.

DIRECTIONS FOR USE

Use as directed. Federal and State laws and regulations may apply. Read and follow all directions on this label.

VEGETABLES

APPLICATION—Apply at the rate of 1/2 to 1 pound per acre. Repeat applications as necessary. Do not apply to plants that are stressed or damaged. Do not apply to plants that are in bloom.

BEANS—To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Leaf Miner, Flea Beetle, Japanese Beetle, and Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Lima Bean Bug, apply at the rate of 1/2 to 1 pound per acre. To control Lima Bean Pod Borer, apply at the rate of 1/2 to 1 pound per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, MUSTARD GREENS, TURNIPS, KOMRABI AND SPINACH—To control Cabbage Root Fly, apply at the rate of 1/2 to 1 pound per acre. To control Cabbage Looper, apply at the rate of 1/2 to 1 pound per acre. To control Imported Cabbageworm, Diamond Back Moth, and Imported Cabbageworm, apply at the rate of 1/2 to 1 pound per acre. To control Cabbage Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Cabbage Aphid, apply at the rate of 1/2 to 1 pound per acre. To control Cabbage Root Borer, apply at the rate of 1/2 to 1 pound per acre.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS—To control Cucumber Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre.

CORNS—To control European Corn Borer, apply at the rate of 1/2 to 1 pound per acre. To control Corn Earworm, apply at the rate of 1/2 to 1 pound per acre. To control Corn Rootworm, apply at the rate of 1/2 to 1 pound per acre. To control Corn Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Corn Aphid, apply at the rate of 1/2 to 1 pound per acre.

FRUIT TREES—To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

ORNAMENTALS—To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

LAWNS—To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

EARWIGS—To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre.

CARROTS

To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

LETTUCE

To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

ONION

To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

POTATO

To control Colorado Potato Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

TOMATO

To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

EGGPLANT

To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

PEPPER

To control Japanese Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Green Earwig, apply at the rate of 1/2 to 1 pound per acre. To control Flea Beetle, apply at the rate of 1/2 to 1 pound per acre. To control Leaf Miner, apply at the rate of 1/2 to 1 pound per acre.

INTERVAL BEFORE HARVEST

See back panel for interval before harvest information.

CAUTION

See back panel for caution information.

DISPOSAL

See back panel for disposal information.

ACTIVE INGREDIENT

75% Cyfluthrin (95% Cyfluthrin, 5% Methoxy Carbamate)

INERT INGREDIENT

25% Inert Ingredients

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE BACK PANEL

NET WT. 2 LB.

RALSTON PURINA COMPANY

GENERAL OFFICE • 860 North American Ave.
St. Louis, Missouri 63164

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Irritation to eyes, nose, throat, and skin. Avoid contact with eyes, nose, throat, and skin. Wash thoroughly with soap and water after handling. Avoid breathing dust or mist.

NOTE FOR PHYSICIAN: See back panel for first aid instructions.

STATEMENT OF PRACTICAL TREATMENT

First Aid: If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If inhaled, get fresh air and rest.

ENVIRONMENTAL HAZARDS: Do not apply to water bodies. Do not apply to areas where birds or bees are present. Do not apply to areas where children or pets play.

DIRECTIONS FOR USE

Read label carefully. Do not use if directions have been changed without your knowledge.

VEGETABLES

APPLICATION: Begin applications when insects first observed and repeat at regular intervals as necessary for good control.

BEANS: To control Mexican Bean Beetle, Flea Beetle, Cucumber Beetle, Leafhopper, Japanese Beetle, and Corn Earworm, apply at the rate of 10 pounds per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, MUSTARD GREENS, TURNIPS, KOHLRABI AND SPINACH: To control Flea Beetle and Harlequin Bug, apply at the rate of 10 pounds per acre.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control Pumpkin Moth, Cucumber Beetle, Squash Bug, Flea beetles, apply at the rate of 10 pounds per acre.

CORN: To control Corn Earworm, European Corn Borer, Armyworm, Saw beetle, and Japanese Beetle, apply at the rate of 10 to 20 pounds per acre.

5112 B

QUALITY CONTROLLED BY PURINA RESEARCH

CARROTS

LETTUCE

To control Flea Beetle, apply at the rate of 10 pounds per acre.

ORRA

TOMATO, EGGPLANT, PEPPER

To control Flea Beetle, apply at the rate of 10 pounds per acre.

POTATO

To control Flea Beetle, apply at the rate of 10 pounds per acre.

INTERVAL BEFORE HARVEST

See back panel for interval before harvest instructions.

CAUTION

Do not apply to areas where children or pets play.

ORNAMENTALS

To control Flea Beetle, apply at the rate of 10 pounds per acre.

LAWNS

To control Flea Beetle, apply at the rate of 10 pounds per acre.

EARWIGS

To control Earwig, apply at the rate of 10 pounds per acre.

DISPOSAL

Do not dispose of this product in water.



PURINA[®]

GENERAL SEVIN[®]-7

INSECTICIDE

Controls Garden and Ornamental Insects

E.P.A. REG. NO. 602

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-methylcarbamate)

NET WEIGHT

9.5
100

Trade Name: 1-naphthyl N-methylcarbamate

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE BACK PANEL

NET WT. 5 LB.

RALSTON PURINA COMPANY

200 North Lincoln Avenue, St. Louis, Missouri 63102

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Harmful if swallowed. Avoid breathing dust or spray. Wash thoroughly after use. Avoid getting eyes washed with water before use. Avoid contact with clothing. Wash hands thoroughly after use.

NOTE FOR PHYSICIAN: Avoid use in pregnant women and nursing infants. Avoid use in children.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, induce vomiting. If inhaled, get fresh air. If on skin, wash with soap and water. If in eyes, flush with water.

ENVIRONMENTAL HAZARDS: Do not apply to water bodies. Do not contaminate water. Do not apply to crops as specified in this label. Bee Caution: This product is highly toxic to bees exposed to direct treatment. Do not apply to crops. Protective clothing should be worn and removed before eating, drinking, or smoking.

DIRECTIONS FOR USE

This application of federal law is subject to the provisions of state law. If state law is inconsistent with this labeling, the state law applies.

VEGETABLES

APPLICATION: Begin application when insect damage is first observed and repeat at regular intervals as necessary for good insect control. Gardeners apply at the rate of 10 to 20 pounds per acre, which is equivalent to 20 pounds per acre.

BEANS: To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Leafhoppers, Flea Beetles, Spanish Beetles, and Corn Earworm, apply at the rate of 10 pounds per acre. To control Legume Bug, the Bug and the Bean Pod Borer, apply 20 pounds per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPROUTS, CALLIFLOWER, MUSTARD GREENS, TURNIPS, KOHLRABI AND SPINACH: To control Flea Beetle and Harlequin Bug, apply at the rate of 10 pounds per acre. To control Armyworm, apply 10 to 20 pounds per acre. Apply 2 days before application and harvest. Broccoli, Brussels Sprouts, Cabbage, Cauliflower and Kohlrabi: Apply 2 days before harvest. Mustard Greens, Spinach and Turnips: Apply 2 days before harvest.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control Pickleworm, Melonworm, Cucumber Beetle, Squash Bug, Flea Beetles, apply at the rate of 10 pounds per acre.

CORN: To control Corn Earworm, European Corn Borer, Armyworm, Sap Beetles, and Japanese Beetles, apply at the rate of 10 to 20 pounds per acre. European Corn Borer apply at the rate of 10 to 20 pounds per acre when plants are about knee high. Repeat at weekly intervals until tassels emerge. For European Corn Borer, apply to the plants and repeat, as necessary, at 3 to 5 day intervals starting when first tassels appear, and continue this schedule until tassels are fully developed. Repeat at weekly intervals, as necessary, until tassels are fully developed. This schedule is for the purpose of reducing the pest population.

CARROTS: To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre. LETTUCE: To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre. OKRA: To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre.

TOMATO, EGGPLANT, PEPPER: To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre.

POTATO: To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre.

INTERVAL BEFORE HARVEST: Apply 2 days before harvest.

CAUTION: Do not apply to water bodies. Do not contaminate water.

ORNAMENTALS

To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre. Do not apply to water bodies. Do not contaminate water.

LAWNS

To control Flea Beetles, apply at the rate of 10 to 20 pounds per acre. Repeat at weekly intervals.

EARWIGS

This material is effective against earwigs. It is used to control earwigs on lawns. Apply at the rate of 10 to 20 pounds per acre. Repeat at weekly intervals.

DISPOSAL: Do not use in water bodies. Do not contaminate water.




PURINA®

GENERAL SEVIN*-7

INSECTICIDE

Controls Garden and Ornamental Insect Pests



ACTIVE INGREDIENT: Gamma-Hydroxy-N-methylcarbamate 93.0%
INERT INGREDIENT: 7.0%

*Trademark of Cyanamid Corp. of Gamma-Hydroxy-N-methylcarbamate

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE BACK PANEL

NET WT. 5 LB.

RALSTON PURINA COMPANY
GENERAL OFFICES • CHECKERBOARD SQUARE
ST. LOUIS 5, MISSOURI 63164

QUALITY CONTROLLED BY PURINA RESEARCH