



EPA Reg. No. 602-324

# RABON DUST

**Contains 3% Rabon\*\* Insecticide****For Use On Livestock, Poultry, Dogs and Cats****No Milk or Tissue Withdrawal • A Ready-To-Use Insecticide Dust**NOT REVIEWED  
In accordance with PR Notice 82-2.  
Based on draft labelling dated 1/13/58

**In Livestock:** Controls Horn Flies, Cattle Grubs and Lice and aids in the control of Face Flies on Beef and Dairy Cattle. Controls Lice on Swine.

**In Poultry:** Controls Northern Fowl Mites, Chicken Mites, and Lice on or around Chickens. Controls Darkling Beetle (litter beetle) in litter.

**In Dogs and Cats:** Controls both Fleas and Ticks. Kills fleas up to 16 days and Brown Dog Ticks up to 7 days. May also be used to help control Fleas and Ticks in pet's sleeping quarters.

ACTIVE INGREDIENT:	By Weight
2-Chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphite	3.00%
INERT INGREDIENTS	97.00%
	100.00%

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Additional Precautionary Statements on Back Panel

### PRECAUCION

Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

### (ENGLISH TRANSLATION) CAUTION

To the User: If you cannot read English, do not use this product until the label has been fully explained to you.

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Flush with plenty of water. If irritation persists, seek medical attention.

**NOTE TO PHYSICIAN/VETERINARIAN:** Rabon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

## PRECAUTIONARY STATEMENTS HAZARD TO HUMANS CAUTION

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash eyes with plenty of water. If irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and food-stuffs. Do not apply in dwellings.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a dry place in original container.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

### WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### FOR USE ON POULTRY

TYPE OF HOUSING	APPLICATION	REMARKS
Wire Cages	1 lb/300 birds Plunger, rotary, type duster or shaker-can duster.	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days. Wire rungs & corners should also be treated.
		<b>NOTE:</b> For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.
Floor Management Litter	1 lb/100 sq. ft. Plunger or rotary type duster.	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	2 lb/100 birds.	Use box about 2 ft. by 2 ft. by 1 ft. deep
Roost Paint	1 lb/100 sq. ft. — make thick slurry by mixing 1 lb. of dust with 1 pint of water, continually stir.	Treat roosts thoroughly, particularly the cracks and crevices.

#### FOR USE ON DOGS AND CATS

To control fleas and ticks on your pet, dust powder liberally over the animal and rub thoroughly through hair coat to skin. Apply also to feet, between the toes and on the legs for tick control. Repeat at weekly intervals if necessary. Occasional dusting of pet's bedding and around pet's sleeping quarters and other areas will help free area of fleas and ticks.

#### FOR USE ON LIVESTOCK

PURINA Rabon Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

Animal	Insect	Remarks
Beef Cattle & Dairy Cattle	Horn Flies & Face Flies & Lice	<b>HAND DUSTING</b> — Apply approximately 2 oz. of dust by shaking can, rotary duster or by spoon to the upper portions of the back as an aid for the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary. <b>SELF-TREATING DUST BAG</b> <b>FORCED USE</b> — Put PURINA Rabon Dust in cotton cloth or double burlap bags or use weather-proof cattle dust bags and hang in door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather. <b>FREE-CHOICE USE</b> — Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.
	Cattle Grubs	After the grubs have encysted (formed warbles) apply 3-4 oz. down the backline and rub in thoroughly, taking care to get the dust into the warble if possible.
Swine	Lice	<b>HAND DUSTING</b> — Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of PURINA Rabon Dust should be applied per 150 sq. ft. of bedding.

\*Rabon is a registered trademark of Shell Oil Company

**QUALITY CONTROLLED BY PURINA RESEARCH**

EPA Est. No. 602-MO-1

Manufactured by

**PURINA MILLS, INC.**

PO Box 66812

St. Louis, Missouri 63166-6812

**NET WT 1 LB**

D-8153

Printed in U.S.A.

ARG/B



# RABON DUST

## Contains 3% Rabon<sup>®</sup>\* Insecticide

### For Use On Livestock, Poultry, Dogs and Cats

### No Milk or Tissue Withdrawal • A Ready-To-Use Insecticide Dust

NOT REVIEWED

In accordance with PR Notice 82-2.

Based on draft labelling dated 1/13/58

**Livestock:** Controls Horn Flies, Cattle Grubs and Lice and aids in the control of Face Flies on Beef and Dairy Cattle. Controls Lice on Swine.

**Poultry:** Controls Northern Fowl Mites, Chicken Mites, and Lice on and Chickens. Controls Darkling Beetle (litter beetle) in litter.

**Dogs and Cats:** Controls both Fleas and Ticks. Kills fleas up to 16 days and Dog Ticks up to 7 days. May also be used to help control Fleas and Ticks in pet's sleeping quarters.

<b>ACTIVE INGREDIENT:</b>	By Weight
2-Chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate	3.00%
<b>INERT INGREDIENTS</b>	97.00%
	100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Additional Precautionary Statements on Back Panel

#### PRECAUCION

Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(ENGLISH TRANSLATION)

#### CAUTION

To the User: If you cannot read English, do not use this product until the label has been fully explained to you.

#### STATEMENT OF PRACTICAL TREATMENT

**TO EYES:** Flush with plenty of water. If irritation persists, seek medical attention.

**NOTE TO PHYSICIAN/VETERINARIAN:** Rabon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS  
CAUTION**

If swallowed: Avoid breathing dust. Do not get in eyes, on skin, or on thing. If the material gets into the eyes, wash eyes with plenty of water. If irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not apply in dwellings.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a dry place in original container.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

#### WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### FOR USE ON POULTRY

TYPE OF HOUSING	APPLICATION	REMARKS
Wire Cages	1 lb./300 birds Plunger, rotary, type duster or shaker-can duster	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days. Wire rungs & corners should also be treated.
		<b>NOTE:</b> For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation or breeding flocks.
Floor Management Litter	1 lb./100 sq. ft. Plunger or rotary type duster	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	2 lb./100 birds	Use box about 2 ft. by 2 ft. by 1 ft. deep.
Roost Paint	1 lb./100 sq. ft. — make thick slurry by mixing 1 lb. of dust with 1 pint of water, continually stir	Treat roosts thoroughly, particularly the cracks and crevices.

#### FOR USE ON DOGS AND CATS

To control fleas and ticks on your pet, dust powder liberally over the animal and rub thoroughly through hair coat to skin. Apply also to feet, between the toes and on the legs for tick control. Repeat at weekly intervals if necessary. Occasional dusting of pet's bedding and around pet's sleeping quarters and other areas will help free area of fleas and ticks.

#### FOR USE ON LIVESTOCK

PURINA Rabon Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

Animal	Insect	Remarks
Beef Cattle & Dairy Cattle	Horn Flies, Face Flies & Lice	<b>HAND DUSTING</b> — Apply approximately 2 oz. of dust by shaking can, rotary duster or by spoon to the upper portions of the back as an aid for the control of face flies. Rub lightly to carry the dust beneath the hair. Repeat as necessary. <b>SELF-TREATING DUST BAG FORCED USE</b> — Put PURINA Rabon Dust in cotton cloth or double burlap bags or use weather-proof cattle dust bags and hang in door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather. <b>FREE-CHOICE USE</b> — Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.
	Cattle Grubs	After the grubs have encysted (formed warbles) apply 3-4 oz. down the backbone and rub in thoroughly, taking care to get the dust into the warble if possible.
Swine	Lice	<b>HAND DUSTING</b> — Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of PURINA Rabon Dust should be applied per 150 sq. ft. of bedding.

\*Rabon Insecticide  
Rabon is a registered trademark of Shell Oil Company

**QUALITY CONTROLLED BY PURINA RESEARCH**

EPA Est. No. 602-MO-1

Manufactured by  
**PURINA MILLS, INC.**  
P.O. Box 66812  
St. Louis, Missouri 63166-6812

**NET WT 1 LB  
D-8153**