

ACCEPTED
 OCT 25
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 602-324

pes per LPA.

DATE ISSUED: 1/21/09 (YELLOW)

PURINA
 BRAND
 Rev. Date Issued: 2/15/88

EPA Reg. No. 602-324

RABON DUST
 Contains 3% Rabon * Insecticide

(3-Row Checkerband®)

For Use On Livestock, Poultry, Dogs and Cats

No Milk or Tissue Withdrawal

A Ready-To-Use Insecticide Dust

On Livestock:

Controls Horn Flies and Lice, and aids in the control of Face Flies on Beef and Dairy Cattle. Controls Lice on Swine.

On Poultry:

Controls Northern Fowl Mites, Chicken Mites, and Lice on or around Chickens. Controls Darkling Beetle (litter beetle) in litter.

On Dogs and Cats:

Controls both Fleas and Ticks. Kills fleas up to 16 days and Brown Dog Ticks up to 7 days. May also be used to help control Fleas and Ticks in pet's sleeping quarters.

ACTIVE INGREDIENT:

By Weight

Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	3.00%
INERT INGREDIENTS	<u>97.00%</u>
	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Additional Precautionary Statements on Back Panel

PRECAUCION

Ai Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(ENGLISH TRANSLATION)

CAUTION

To the User: If you cannot read English, do not use this product until the label has been fully explained to you.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. If irritation persists, seek medical attention.

NOTE TO PHYSICIAN/VETERINARIAN: Rabon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

NET WT 1 LB

D-8153

Manufactured By
 PURINA MILLS, INC.
 P. O. Box 66812
 St. Louis, Missouri 63166-6812 U.S.A.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS
CAUTION**

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash eyes with plenty of water. If irritation persists, see a physician. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not apply in dwellings.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff may be hazardous to aquatic organisms in adjacent areas. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwater. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


Storage: Store in a dry place in original container.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this Warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


FOR USE ON POULTRY

TYPE OF HOUSING	APPLICATION	REMARKS
Wire Cages	1 lb/300 birds Plunger, rotary, type duster or shaker-can duster.	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days. Wire rungs & corners should also be treated.
NOTE: For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.		
Floor Management Litter	1 lb/100 sq. ft. Plunger or rotary type duster.	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	2 lb/100 birds.	Use box about 2 ft. by 2 ft. by 1 ft. deep.
Roost Paint	1 lb/100 sq. ft. --make thick slurry by mixing 1 lb. of dust with 1 pint of water, continually stir.	Treat roosts thoroughly, particularly th. cracks and crevices.

FOR USE ON DOGS AND CATS

To control fleas and ticks on your pet, dust powder liberally over the animal and rub thoroughly through hair coat to skin. Apply also to feet, between the toes and on the legs for tick control. Repeat at weekly intervals if necessary.

Occasional dusting of pet's bedding and around pet's sleeping quarters and other areas will help free area of fleas and ticks.



ACCEPTED
 OCT 25 1988
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 602-324

rev. Date Issued: 2/15/88

**PURINA®
 BRAND**

EPA Reg. No. 602-324

RABON DUST
 Contains 3% Rabon * Insecticide

(3-Row Checkerband®)

For Use On Livestock, Poultry, Dogs and Cats

No Milk or Tissue Withdrawal

A Ready-To-Use Insecticide Dust

On Livestock:

Controls Horn Flies and Lice, and aids in the control of Face Flies on Beef and Dairy Cattle. Controls Lice on Swine.

On Poultry:

Controls Northern Fowl Mites, Chicken Mites, and Lice on or around Chickens. Controls Darkling Beetle (litter beetle) in litter.

On Dogs and Cats:

Controls both Fleas and Ticks. Kills fleas up to 16 days and Brown Dog Ticks up to 7 days. May also be used to help control Fleas and Ticks in pet's sleeping quarters.

ACTIVE INGREDIENT:

By Weight

Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate 3.00%

INERT INGREDIENTS..... 97.00%
 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Additional Precautionary Statements on Back Panel

PRECAUCION

Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
 (ENGLISH TRANSLATION)

CAUTION

To the User: If you cannot read English, do not use this product until the label has been fully explained to you.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. If irritation persists, seek medical attention.

NOTE TO PHYSICIAN/VETERINARIAN: Rabon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

NET WT 1 LB

D-8153

Manufactured By
 PURINA MILLS, INC.
 P. O. Box 66812
 St. Louis, Missouri 63166-6812 U.S.A.

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS
CAUTION**

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash eyes with plenty of water. If irritation persists, see a physician. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not apply in dwellings.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff may be hazardous to aquatic organisms in adjacent areas. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwater. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE ON POULTRY

TYPE OF HOUSING	APPLICATION	REMARKS
Wire Cages	1 lb/300 birds Plunger, rotary, type duster or shaker-can duster.	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days. <u>Wire rings & corners should also be treated.</u>
NOTE: For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.		
Floor Management Litter	1 lb/100 sq. ft. Plunger or rotary type duster.	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	2 lb/100 birds.	Use box about 2 ft. by 2 ft. by 1 ft. deep.
Roost Paint	1 lb/100 sq. ft. --make thick slurry by mixing 1 lb. of dust with 1 pint of water, continually stir.	Treat roosts thoroughly, particularly the cracks and crevices.

FOR USE ON DOGS AND CATS

To control fleas and ticks on your pet, dust powder liberally over the animal and rub thoroughly through hair coat to skin. Apply also to feet, between the toes and on the legs for tick control. Repeat at weekly intervals if necessary.

Occasional dusting of pet's bedding and around pet's sleeping quarters and other areas will help free area of fleas and ticks.

FOR USE ON LIVESTOCK

PURINA Rabon Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

<u>Animal</u>	<u>Insect</u>	<u>Remarks</u>
Beef Cattle & Dairy Cattle	Horn Flies Face Flies & Lice	<p>HAND DUSTING - Apply approximately 2 oz. of dust by shaking can, rotary duster or by spoon to the upper portions of the back as an aid for the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.</p> <p>SELF-TREATING DUST BAG</p> <p>FORCED USE - Put PURINA Rabon Dust in cotton cloth or double burlap bags or use weather-proof cattle dust bags and hang in door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.</p> <p>FREE-CHOICE USE - Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.</p>
Swine	Lice	<p>HAND DUSTING - Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of PURINA Rabon Dust should be applied per 150 sq. ft. of bedding.</p>

QUALITY CONTROLLED BY PURINA RESEARCH

*Rabon Insecticide

Rabon is a registered trademark of Shell Oil Company

8901 D

Printed in U.S.A.