

Pm 10

602-330

Rev. Date Issued: 6/24/85

(9-Square Logo )

PURINA

(3-Row Checkerband)

FLY-A-REST AQ BRAND

PROFESSIONAL STRENGTH - LOW DOSAGE REQUIRED

FOR USE IN SCHOOLS, HOSPITALS, INSTITUTIONS, SUPERMARKETS, DAIRIES, HOTELS, RESTAURANTS, DRIVE-INS, MOTELS, DRUG STORES, FOOD PLANTS, MILK ROOMS AND ANIMAL QUARTERS.

PROVIDES RAPID CONTROL OF HOUSE FLIES, MOSQUITOES, SMALL FLYING MOTHS (MILLERS), GNATS, WASPS AND ACCESSIBLE STAGES OF COCKROACHES, ANTS, WATER BUGS, SILVERFISH, FLOUR BEETLES, WEEVILS, GRAIN MITES AND SPIDERS.

KILLS AND REPELS STABLE FLIES, HORSE FLIES, FACE FLIES, DEER FLIES, HOUSE FLIES, HORN FLIES, MOSQUITOES AND GNATS.

ACTIVE INGREDIENTS:

Pyrethrins..... 0.5%
\*Piperonyl Butoxide, Technical ..... 5.0%

INERT INGREDIENTS..... 94.5%
100.0%

\* Equivalent to 4% (butylcarbityl) (6-propylpiperonyl) ether and 1% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED - Remove affected person to fresh air. Apply artificial respiration if indicated.


IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES - Flush eyes with plenty of water. Call a physician if irritation persists.

D-

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812
St. Louis, Missouri 63166-6812 U.S.A.

EPA Est. No. 602-MO-1
EPA Reg. No. 602-



NET WT 24 OZ

602-330

**Right Side Panel****PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds, and cover fish aquariums before spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of waste or cleaning of equipment.

**PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° may cause bursting.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry area away from heat or open flame.

**PESTICIDE DISPOSAL:** Securely wrap container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Replace cap and discard container in trash. Do not incinerate or puncture.

**SHAKE BEFORE USING**

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray in close range.

**House flies, mosquitoes, gnats, wasps and small flying moths (Millers):** Close all doors and windows and spray room at the rate of 2 seconds per 1000 cubic feet. Direct spray to all parts of the room especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10-15 minutes after treatment. Repeat as necessary.

**Cockroaches, waterbugs, silverfish, crickets, spiders, centipedes and sow bugs:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

#### Left Side Panel

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**Fleas:** Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

**Carpet Beetles:** Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.


**Bedbugs:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

**In Animal Quarter and Milk Rooms:** For rapid control of house flies, mosquitoes, gnats, wasps and small flying moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

**To Treat Animals:** To protect beef and dairy cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

**Exposed Stages Of Indian Moths And Angoumois Grain Moths:** Close doors and windows. Directly spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

**Exposed Stages Of Flour Beetles, Weevils And Grain Mites:** Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.



**Controls Garden Insects on Ornamental and Vegetable Plants:** Spray with slow sweeping motion, not closer than 18 inches from plants. Apply in accordance with the chart below. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts for 1-2 seconds. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application of 7 to 10 day intervals or after heavy rains.

Acceptable for use to day of harvest on edible plants.

### APPLICATION

<u>USE ON</u>	<u>SECONDS OF SPRAY/ FOOT OF ROW OR PER PLANT</u>	<u>TO CONTROL</u>
Roses	2-3	Aphids, rose chafers, Japanese beetles, rose slugs, tent caterpillars, scales.
Asters	2-3	Leafhoppers, tarnish plant bugs, lace bugs, thrips (exposed) Mealy Bugs.
Dahlias	2-3	Aphids, red spider mites, plant bugs
African Violets	1	Broad mites, citrus mealybugs.
Azaleas	1-2	Azalea lace bugs, white flies, leaf miners.
Euonymus	1-2	Euonymus scales, aphids, thrips (exposed).
Hydrangeas	1-2	Aphids red spider mites, hydrangea leaf tiers.
Junipers	3-5	Juniper scales, red spider mites, bagworms.
Lilacs	3-5	Leaf miners, leaf rollers scales.
Tomatoes	2-4	Flea beetles, hornworms, stink bugs, Colorado potato beetles, green peach aphids.
Peppers, Eggplants Potatoes	1-2	Colorado potato beetles, leafhoppers, blister beetles, flea beetles, green peach aphids.
Cucumbers, Squash Melons	2-4	Striped and spotted cucumber beetles.
Beans	1-2	Mexican bean beetles, white flies, 12-spotted beetles.
Swee. Corn	2-3	Japanese beetles.