



Purina

HARD HITTER[®] Brand CATTLE POUR-ON INSECTICIDE

Controls Lice and Flies on Dairy and Beef Cattle



ACTIVE INGREDIENTS

Permethrin (3-Phenoxyphenyl) methyl (±) cis, trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate*
INERT INGREDIENTS

Contains petroleum distillates

1.0%
99.0%
100.0%

*Cis/trans ratio: Min. 35% (±) cis and max. 65% (±) trans


ACCEPTED
JUN - 4 1954
U.S. Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the purpose of registration
EPA Reg. No. 602-331

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

Manufactured By
PURINA MILLS, INC.
P.O. Box 66812
St. Louis, Missouri 63166-6812

EPA Reg. No. 602-331

NET 1 GAL



PURINA[®] HARD HITTER[®] CATTLE POUR-ON INSECTICIDE Brand

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with eyes.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

If swallowed: Call a physician immediately. Do not induce vomiting unless under medical attention.

NOTE TO PHYSICIAN

Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by disposal of equipment washwaters. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE—No dilution necessary.

Apply to Lactating and non-lactating dairy cattle, beef cattle and calves.

APPLICATION

TARGET INSECTS APPLICATION INSTRUCTIONS

Lice
Horn flies
Face flies

Beginning at the head, pour along the center of the back. Apply 1/2 fluid ounces per 100 pounds body weight of animal up to a maximum of 5 fluid

Repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control, a double treatment at 14-day intervals is recommended.

Special Note: PURINA HARD HITTER Cattle Pour-On Insecticide is not effective in controlling cattle grubs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.