Reason For Issue: Reg # 602 - 358

Adding rats to label.

PM-14 Date Issued: 3/19/93 (Green) Prev. Date Issued: None UPC Code: 01929801316

FRONT PANEL

(9-Square Logo')

EPA Reg. No. 602-

PURINA '

ASSAULTTM MOUSE/RAT

PLACE PACK [·] BRAND JAN 0 5 1994 Under the Federal Insecticide, Fungistice, and Redenticide Act as assended, for the particide replaced under EPA Reg. No. 602 - 358

ACCEPTED

Mouse and Rat Poison For Indoor and Outdoor Rodent Control

Kills House Mice, Norway Rats and Roof Rats

House Mice, Norway Rats and Roof Rats usually consume a lethal dose in a single day's feeding, but it may take two or more days from the time of bait consumption, or pack removal, for these rodents to die. Rodents cease feeding after consuming a lethal dose. ASSAULT Mouse/Rat Place Pack is effective against anticoagulant-resistant mice and rats.

| INERT INGREDIENTS | <u>99.99%</u> |
|-------------------|---------------|
| TOTAL | 100.00% |

| KEEP OUT OF REACH OF CHILDREN |
|-------------------------------|
| CAUTION |
| SEE OTHER CAUTIONS BELOW |
| |

NET WT 25 x 0.5 Oz

D-0212

Manufactured By PURINA MILLS, INC. P.O. Box 66812 St. Louis, Missouri 63166-6812 U.S.A.

CONTAINS BITREX

Right Panel

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed. Avoid contact with skin, eyes or clothing. In case of accidental contact, flush eyes or skin with water and wash contaminated clothing. Seek medical attention immediately if ingested or if breathing difficulty occurs or irritation persists. Wash hands thoroughly after handling Assault Mouse/Rat Place Pack.

NOTE TO PHYSICIAN:

Assault Mouse/Rat Place Pack is not an anticoagulant type of rodenticide. If ingested, limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamperresistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
 - 3. Dispose of preduct container and unused, spoiled, and unconsumed bait as specified to this label.

USE RESTRICTIONS - For control of Norway rats and roof rats in and around homes, industrial and agricultural buildings, and similar man-made structures. This product may also be used in alleys located in unban areas, inside transport vehicles (ships, trains, aircraft), and in and around related port or terminal buildings. Do not use ASSAULT MOUSE/RAT PLACE PACK in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS - Determine areas where rats will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

The active ingredient in ASSAULT MOUSE/RAT PLACE PACK is different from anticoagulant rodenticides. Generally, rodents cease feeding after consuming a lethal amount and die within a few days.

RATS: Apply 2 - 12 place packs per placement location. Space placements at intervals of 15 to 30 feed.

MICE: Apply 1 - 2 place packs per placement. Space placements at intervals of 8 to 12 feet. Larger placements. up to six place packs, may be needed at points of extremely high mouse activity.

RATS and MICE: Adjust the amount of bait applied to the level of feeding by target rodents expected at each placement location. Highest bait consumption is expected to occur during the first day or two after treatment. After several days, inspect bait placements and replenish bait at sites where there is heavy feeding or where there is continued evidence of rodent activity. Maintain bait for at least one week and continue baiting until all signs of rodent activity have ceased or it becomes clear that the remaining rodents are not attracted to the bait.

Collect and properly dispose of all dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.


STORAGE AND DISPOSAL

Do not contamine water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry place away from other chemicals. Keep containers closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty containers. Dispose of liner in a sanitary landfill or by incineration if allowed by state or local authorities. If burned, stay out of smoke.



9303 A