

604-24

hammond
SINCE 1875



slug shot
garden dust

ACCEPTED
5-1-37
604-24
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MOLLUSCIDICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 10000 SUBJECT
TO ATTACHED COMMENTS.

SLUG SHOT DUST

NET WEIGHT 1 LB.

DIRECTIONS FOR USE

The following general directions are supplied by us gratuitously and are believed to be reliable, but are in no way guaranteed. Agricultural Experiment Stations in many States issue recommendations for use of Rotenone under local conditions.

Application: Apply carefully with a good duster so as to obtain a light, even coverage on both the upper and lower sides of the leaves. Applications should be repeated after heavy rains. Dusting will be most effective when done when the air is still. ~~Do not apply to food crops within one day of harvest. Keep out of reach of children.~~

USES

VEGETABLES

On Asparagus: For Asparagus Beetles apply when insects first appear, repeat in 5 to 7 days if needed.

On Beans: For Mexican Bean Beetle apply to the underside of the leaves when eggs begin to hatch. Repeat applications at 7 day intervals if needed. For Bean Leaf Beetle apply when insects are first seen, repeat at 7 day intervals if needed.

On Cabbage, Cauliflower, Broccoli and Other Crucifers: For Cabbage Worms begin applications as soon as plants are established in the field, repeat at 7 day intervals if needed.

On Melons, Cucumber, Squash and Other Cucurbits: For striped and Spotted Cucumber Beetles and Aphids, begin applications when insects first appear and repeat at 7 day intervals if needed.

On Peas: For Aphids and Weevils apply when insects first appear and repeat once or twice at 5 to 7 day intervals if needed. Under conditions of severe weevil infestation a fourth application may be needed.

On Sweet Corn: For European Corn Borer make first application when eggs are hatching and make 3 additional applications at 5 day intervals. Direct the dust downward into the whorls.

~~Residues of this product should be removed from edible parts of food crops at harvest.~~

FLOWERS

On Calendula, Carnation, Chrysanthemum, Dahlia, Marigold, Nasturtium, Rose, Stock and Sweet Pea: For thrips, caterpillars, and many Aphids, apply when insects are present. Repeat applications at 5 to 7 day intervals or as needed.

SLUG SHOT non-injurious Insecticide Used Continuously for Three Quarters Century	
Active Ingredients:	
Rotenone	.75%
Other Cube Resins	.75%
Sulphur (300 Mesh)	8.00%
Copper (as metallic form Basic Copper Sulphate)	.50%
Inert Ingredients	90.00%

A favorite in home gardens for control of many insect pests on flowers and vegetables.

NOTICE

Seller makes no warranty of any kind, express or implied, concerning the use of this product. The user assumes all risks for use or handling, whether or not in accordance with any direction or suggestion of the seller or its employees.

HAMMOND PAINT & CHEMICAL CO. INC., BOSTON, MASS.