CAPTAN FP 700 R is a microfine wettable powder especially formulated for the treatment of seed, prior to storage and planting, to protect seed from molds and other funglicausing storage loss, and to protect seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

COMPATIBILITY: CAPTAN FP 700 R can be used in combination with most commonly used seed protectants, both insecticides and fungicides as well as fertilizers. Do not use with strongly alkaline materials.

PLANT SAFETY: At the recommended dosage CAPTAN FP 700 R is non-injurious from the standpoint of seed injury or yield reduction, the germination of treated seed is frequently stimulated giving a superior result as compared to results from other seed fungicides.

GENERAL USE PRECAUTIONS

Read all directions before using and use only as specified on this label.

Toxic to fish, keep out of lakes, streams or ponds, do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of excess treated seeds by burying with wastes.

IMPORTANT: Treated seed must not be used for human consumption nor as a feed. Bags containing treated seed should be labeled:

TREATED SEED

DO NOT USE FOR FEED, FOOD OR OIL PURPOSES.

The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

DIRECTIONS FOR USE

SLURRY SEED TREATMENT: Pre-mix the recommended amount of this material with a little water in a bucket and make a thin paste, stir this paste into the required amount of water for the slurry equipment to be used.

NOTE: Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas. If seed is bagged when wet, heat is developed and impairs seed germination.

CONDITIONS OF SALE

- I. GUARD CHEMICAL COMPANY warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
- 2. Critical and unforseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but ere not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above), resulting from handling, storage, and use of this material.

CAPTAN FP-700R

A FUNCICIDE FOR PLANT DIEASE CONTROL

EPA REG. NO. 615-39

CAUTION

ACC PORTON REGISTER KEEP OUT OF REACH OFCHILDREN

MAY BE HARMUL IF SWALLOWED OR INHALED.

AVOID CONTACT WITH SKIN, EYE OR CLOTHING.

AVOID BREATHING DUST, USE WITH ADEQUATE VENTILATION.

WEAR RESPIRATOR, GOGGLES AND GLOVES,

WASH THOROUGHLY AFTER HANDLING.

DO NOT STORE OR USE NEAR FOOD OR FEED.

FIRST AID: IN CASE OF CONTACT IMMEDIATELY FLUSH EYES WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES. CALL A PHYSICIAN. WASH SKIN WITH SOAP AND WATER.

PROTECT FROM EXCESSIVE HEAT -- STORE IN A COOL DRY PLACE.

CONTAINER SHOULD BE DEPOSED BY CRUSHING AND BURYING WITH WASTES.

This product is toxic to fish. Keep out of any body of water.

Do not apply when weather conditions favor drift : f spray from areas treated.

See other panels for directions for use and dosages.

. LES., NET

Manufactured by:

Newark, New Jersey 07105

THE FOLLOWING TABLE gives 4

SEED

CORN-Field

CORN—Sweet

WHEAT For Seed and Seedling Rots. Not

for Bunt

Smut

FLAX

BARLEY

BEANS (Except Baby Limes),
PEAS,
CUCUMBERS
CANTALOUPE

SOUASH
PUMPKIN
WATERMELON
BEETS,
SWISS CHARD
SUGAR BEETS
CLOYER
ALFALFA
SOYBEANS

COTTON—
Acid Delinted
COTTON—
Machine Delinted
Fuzzy (not
delinted)

BLUE GRASS

CABBAGE
CAULIFLOWER,
BROCCOLI,
BRUSSELS
SPROUTS

SPINACH
SORGHUM

hulled (Kernel Smut)
Texas Only
PEPPERS
(Calif. Wonder)
COWPEAS

RICE RYE SESAME especially formulated for the protect seed from molds and from seed-borne

mbination with most commonly as well as fertilizers. Do not

FP 700 R is non-injurious from germination of treated seed is area to results from other seed

NS

ified on this label.

do not contaminate water by

5.

an consumption nor as animal GIC DE

L PURPOSES.

regulations for the application ing treated seed.

amount of this material with a required

be bagged until such time as all ventilated areas. If seed is germination.

naterial conforms to the chemiuse as directed hereon. Manurepresentative to make, any juarantee or representation, ex-

rufacturer's control prevent it f chemicals. Such risks include, which the material is applied, to other plants or crops. Such for the uses stated hereon and user acknowledge and assume manufacturer under 1 above), rial.

GALLOTOX CAPTAN FP-700R

A FUNCICIDE FOR PLANT DISEASE CONTROL

EPA REG. NO. 605-39

CAUTION

TOR ELONGING POISON REGISTERKEEP OUT OF REACH OF CHILDREN

MAY BE HARMUL IF SWALLOWED OR INHALED.

AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

AVOID BREATHING DUST, USE WITH ADEQUATE VENTILATION.

WEAR RESPIRATOR, GOGGLES AND GLOVES,

WASH THOROUGHLY AFTER HANDLING.

DO NOT STORE OR USE NEAR FOOD OR FEED.

FIRST AID: IN CASE OF CONTACT IMMEDIATELY FLUSH EYES WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES. CALL A PHYSICIAN. WASH SKIN WITH SOAP AND WATER.

PROTECT FROM EXCESSIVE HEAT --- STORE IN A COOL DRY PLACE.

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES.

This product is toxic to fish. Keep out of any body of water.

Do not apply when weather conditions favor drift of spray from areas treated.

See other panels for direction for use and dosages.

..... LBS., NET

Manufactured by:

GUARD CHEMICA. COMPANY Newark, New Jersey 07105 THE FOLLOWING TABLE gives amount of CAPTAN FP-700 R to use on various seeds.

SEED	Dosage Rate	Lbs. Seed Treated/lb. of Product
CORN—Field	11/4 oz/100 lbs. 2/3 oz/bu	1280
.CORN-Sweet	2 ² / ₃ oz/100 lbs. 1 ¹ / ₂ oz/bu	600
WHEAT For Seed	11/4 oz/100 lbs. 2/3 oz/bu	1280
OATS Seedling Rots. Not for Bunt	2 oz/100 lbs. 2/3 oz/bu	800
Smut	1½ oz/100 lbs. 3/3 oz/bu 2½ oz/100 lbs.	1070
BEANS (Except Baby Limas),	1½ oz/bu 1¾ oz/100 lbs.	910
PEAS, CUCUMBERS	1 oz/bu 2 oz/100 lbs.	800
SQUASH	11/3 oz/100 lbs.	1200
PUMPKIN_ WATERMELON_		
SWISS CHARD	8 oz/100 lbs.	
SUGAR BEETS	8 oz/100 lbs.	200
CLOVER ALFALFA	5½ oz/100 lbs. 3⅓ oz/bu	290
SOYBEANS	12/3 oz/100 lbs. 1 oz/bu 2 oz/100 lbs.	960
COTTON— Acid Dalinted COTTON—	3 oz/100 lbs.	800
Machine Delinted Fuzzy (not delinted)		530
BLUE GRASS	5½ oz/100 lbs.	290
CABBAGE CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS	1 oz/100 lbs.	1600
SPINACH	4 oz/100 lbs.	400
hulled (Kernel Smut)	3 oz/100 lbs.	530
Texas Only	2 _oz/100 lbs.	800
(Calif. Wonder)	2 oz/100 lbs.	800
COWPEAS	2 67/100 lbs. 11/4 02/5u	800
RICE	21/4 oz/100 lbs.	710
_RYESESAME	13/4 oz/100 lbs.	910
JEJAME	1 oz/100 bs.	1600