

Corn King

DRY INSECTICIDE

Pre-mixed Treatment - For control of chicken red mite, northern fowl mite and poultry lice. Apply 1 lb. per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas. Bird Treatment - Dust powder thoroughly into bird's feathers.

CAUTION: Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid contamination of food and water troughs. Do not reuse container. Destroy container when empty. Do not contaminate water by cleaning equipment, machinery, waste and containers.

ACCEPTED

H. G. Bennett



DRY INSECTICIDE

A POWDERED INSECTICIDE FOR USE ON LIVESTOCK AND POULTRY

Active Ingredients	
Malathion*	4.00%
Inert Ingredients	96.00%

*O,O-dimethyl phosphorid thiocarbonyl diethyl mercaptosuccinate

Caution: Keep out of reach of children. See other warnings on side panel.

25 POUNDS

NET WEIGHT

Manufactured by
THE CORN KING Division
KING CASTLE, Inc.
 Marion, Iowa 52302

Corn King

DRY INSECTICIDE

DIRECTIONS FOR USE

DAIRY CATTLE: Horn Flies — Apply 2 heaping tablespoonsful (2 oz.) to back and neck of each animal. Rub powder into hair. Repeat application at 10-14 day intervals if needed. **Do not apply to lactating dairy animals within 5 hours of milking or during milking.**

BEEF CATTLE: — Lice, Ticks and Horn Flies—Make thorough and liberal application to all parts of the animal's body by hand or power duster. Rub in lightly to carry powder beneath the hair. Repeat only as necessary.

HOGS: — Lice — Make thorough and liberal application to all parts of the animal by hand or power duster as well as pen and litter. Repeat in 10 days if necessary.

SHEEP: — Lice and Keds — Make thorough and liberal application to all parts of the animal by hand or power duster. Repeat only as necessary.

Corn King

DRY INSECTICIDE

POULTRY: Lice and Mites

Apply to the neck, under the wings, and in the tail feathers. Repeat every 7 to 10 days. Do not apply to birds that are stressed or ill.

CAUTION:

Keep out of reach of children. Do not use on humans. Avoid contact with eyes, nose, and mouth. Wash hands thoroughly after use.



DRY INSECTICIDE

A POWDERED INSECTICIDE FOR USE ON LIVESTOCK AND POULTRY

Active Ingredient: Malathion 4.00
Inert Ingredients 96.00

10,0-dimethyl phosphorothioate of diethyl mercaptosuccinate

Caution: Keep out of reach of children. See other warnings on side panel.

25 POUNDS

NET WEIGHT

Manufactured by
THE CORN KING Division
KING CASTLE, Inc.
Marion, Iowa 52302

Corn King

DRY INSECTICIDE

DAIRY CATTLE: Horn Flies — Apply to the face, neck, and body. Repeat every 7 to 10 days. Do not apply to lactating dairy animals within 5 hours of milking or during milking.

BEEF CATTLE: — Lice, Ticks and Horn Flies — Apply to the neck, under the wings, and in the tail feathers. Repeat every 7 to 10 days.

HOGS: — Lice — Apply to the neck, under the wings, and in the tail feathers. Repeat every 7 to 10 days.

SHEEP: Lice and Keds

Apply to the neck, under the wings, and in the tail feathers. Repeat every 7 to 10 days.