

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION

Harmful if inhaled or swallowed. Avoid breathing spray or mist. Do not take internally. Avoid contact with skin, eyes or clothing. In case of contact and before eating and smoking, wash thoroughly with warm water and soap.

Note to Physician: Malathion and Carbaryl are moderate, reversible Cholinesterase inhibitors. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage: Keep container tightly closed to prevent evaporation. Do not store near or contaminate feed or foodstuffs. Do not use, pour, spill or store near heat or open flame. Store in a cool, dry area inaccessible to children and pets.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

BEST AVAILABLE COPY

Corn King

DRY INSECTICIDE

A POWDERED INSECTICIDE FOR USE ON LIVESTOCK AND POULTRY.

Active Ingredients:

Malathion*	4.00%
Inert Ingredients	96.00%

*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water immediately and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air and apply respiration if indicated.

If on skin - Wash promptly with soap and water. Rinse thoroughly.

If in eyes - Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: 2-PAM is antidotal and may be administered in conjunction with atropine sulfate.

25 POUNDS

SEE SIDE PANELS FOR ADDITIONAL
DIRECTIONS FOR USE

NET WEIGHT

Lot No. _____

Avoid
using

HOGS
the a
neces
BEEF
appi
light
DAIR
back
10-14
hours
SHEE
of the
POUL
red m
of litt
ment
CAU
thoro
not re
water
on an
E P.A.

PF
A

Under
Rodeo
pact
L.A.R.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
CAUTION**

Avoid breathing of dust. Avoid contact with skin, wash thoroughly after using. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (e.g., swamps, bays, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on flowering crops or weeds. Do not apply this product if there are bees flying around the treatment area.

DIRECTIONS FOR USE

HOGS: — Hogs, Lice — Make thorough and liberal application to all parts of the animal by hand or power duster as well as pen and litter. Repeat only as necessary.

BEEF CATTLE: — Lice, Ticks and Horn Flies — Make thorough and liberal application to all parts of the animal's body by hand or power duster. Rub in lightly to carry powder beneath the hair. Repeat only as necessary.

DAIRY CATTLE: — Horn Flies — Apply 2 heaping tablespoonsful (2 oz.) to back and neck of each animal. Rub powder into hair. Repeat application at 10-14 day intervals if needed. Do not apply to lactating dairy animals within 5 hours of milking or during milking.

SHEEP: — Lice and Keds — Make thorough and liberal application of all parts of the animal by hand or power duster. Repeat only as necessary.

POULTRY: — Lice and Mites — Premise Treatment - For control of chicken red mite, northern fowl mite and poultry lice, apply 1 lb per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas. Bird Treatment - Dust powder thoroughly into bird's feathers.

CAUTION: Avoid breathing of dust. Avoid contact with skin, wash thoroughly after using. Avoid contamination of feed and water troughs. Do not re-use container. Destroy container when empty. Do not contaminate water by cleaning equipment or disposal of waste and containers. Do not use on animals under one month of age.

E.P.A. Reg. No. 606-23

E.P.A. Est. No. 606-IL-1

**Manufactured by
PRINCE AGRI PRODUCTS, INC.**

**ACCEPT ONE PRINCE PLAZA
QUINCY, IL 62301**

JAN 30 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide registered under EPA Reg. No. 606-23

Handwritten mark: 1/2