

DIRECTIONS

HORSES: Lice and Ticks — Use 1 pint in 40 gals. of water. Mange (Sarcoptic and Psoroptic) — Use 1 pint in 20 gals. of water. Apply sufficient diluted spray to assure complete coverage. This may be from 1/2 to 1 gal. depending on the size of the animal. Repeat only as necessary.

RESIDUAL SPRAY: For control of Flies, Gnats, Mosquitoes, Ants and Roaches — For use in farm buildings. Mix 1 pint in 4 gals. of water and spray walls, ceilings, window sills and wherever insects congregate. Cover feed and water troughs. Apply to point of run-off, usually 1 gal. per 500 sq. ft. of surface area. Remove animals from area while spraying. Repeat as necessary.

CAUTION: May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin. Harmful if swallowed. In case of contact wash immediately with soap and water. Avoid contamination of feed and food stuffs. Do not use on household pets or humans. Do not reuse empty containers. Destroy when empty. Use only the 1 pt. to 40 gal. dilution on young farm animals. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not spray animals within 30 days of slaughter or dip within 60 days of slaughter. Do not use on dairy animals. Do not use in poultry houses or dairy barns.

COMBUSTIBLE LIQUID: DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. KEEP CONTAINER CLOSED.



LIVESTOCK MANGE & LICE LIQUID

A LINDANE CONCENTRATE TO BE DILUTED WITH WATER FOR APPLICATION ON FARM ANIMALS AND WITHIN FARM BUILDINGS.

Active Ingredients:	
Gamma Isomer of Benzene Hexachloride (from Lindane)	10.00%
Xylene	85.00%
Inert Ingredients	5.00%

Caution: Keep out of reach of children. See other warnings on side panel.

CONTENTS 1 PINT

THE CORN KING Division
KING CASTLE, Inc.
Marion, Iowa 52302

DIRECTIONS

Use as a spray or dip diluted with water as follows:

SWINE: Lice — 1 pint in 40 gals. of water. Mange (Sarcoptic) — 1 pint in 20 gals. of water. Be sure of complete coverage allowing approximately one quart of diluted spray per adult animal. Repeat only as necessary. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

CATTLE: Beef — Lice and Ticks — Use 1 pint in 40 gals. of water. Mange (Sarcoptic and Psoroptic) — Use 1 pint in 20 gals. of water. Allow approximately 2 qts. of diluted spray per adult animal. Repeat only as necessary.

SHEEP and GOATS: Lice and Keds (Sheep Tick) — Use 1 pint in 40 gals. of water; Scabies (Sheep Mange) — Use 1 pint in 20 gals. of water. Be sure of complete coverage with penetration down to the skin of animal. Allow from 1 to 2 quarts of diluted spray per adult animal depending on wool coat. Treatment right after shearing is most desirable. Repeat only as necessary. Do not use on dairy goats. See left label panel CAUTION for accepted dosage on young livestock; also other necessary livestock cautions. Do not treat animals under 3 months of age.

U.S.D.A. REG. NO. 606 79

MADE IN U.S.A.

ACCEPTED WITH COMMENTS

ACCEPTED

Jan. 8, 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 606-79

Comments

DIRECTIONS

HORSES: Lice and Ticks — Use 1 pint in 40 gals. of water. Mange (Sarcoptic and Psoroptic) — Use 1 pint in 20 gals. of water. Apply sufficient diluted spray to assure complete coverage. This may be from 1/2 to 1 gal. depending on the size of the animal. Repeat only as necessary.

RESIDUAL SPRAY: For control of Flies, Gnats, Mosquitoes, Ants and Roaches — For use in farm buildings. Mix 1 pint in 4 gals. of water and spray walls, ceilings, window sills and wherever insects congregate. Cover feed and water troughs. Apply to point of run-off, usually 1 gal. per 500 sq. ft. of surface area. Remove animals from area while spraying. Repeat as necessary.

CAUTION: May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin. Harmful if swallowed. In case of contact wash immediately with soap and water. Avoid contamination of feed and food stuffs. Do not use on household pets or humans. Do not reuse empty containers. Destroy when empty. Use only the 1 pt. to 40 gal. dilution on young farm animals. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not spray animals within 30 days of slaughter or dip within 60 days of slaughter. Do not use on dairy animals. Do not use in poultry houses or dairy barns.

COMBUSTIBLE LIQUID: DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. KEEP CONTAINER CLOSED.



LIVESTOCK MANGE & LICE LIQUID

A LINDANE CONCENTRATE TO BE DILUTED WITH WATER FOR APPLICATION ON FARM ANIMALS AND WITHIN FARM BUILDINGS.

Active Ingredients:

Gamma Isomer of Benzene Hexachloride (from Lindane)	10.00%
Xylene	85.00%
Inert Ingredients	5.00%

Caution: Keep out of reach of children. See other warnings on side panel.

CONTENTS

1 PINT

**THE CORN KING Division
KING CASTLE, Inc.
Marion, Iowa 52302**

DIRECTIONS

Use as a spray or dip diluted with water as follows:

SWINE: Lice — 1 pint in 40 gals. of water. Mange (Sarcoptic) — 1 pint in 20 gals. of water. Be sure of complete coverage allowing approximately one quart of diluted spray per adult animal. Repeat only as necessary. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

CATTLE: Beef — Lice and Ticks — Use 1 pint in 40 gals. of water. Mange (Sarcoptic and Psoroptic) — Use 1 pint in 20 gals. of water. Allow approximately 2 qts. of diluted spray per adult animal. Repeat only as necessary.

SHEEP and GOATS: Lice and Keds (Sheep Tick) — Use 1 pint in 40 gals. of water; Scabies (Sheep Mange) — Use 1 pint in 20 gals. of water. Be sure of complete coverage with penetration down to the skin of animal. Allow from 1 to 2 quarts of diluted spray per adult animal depending on wool coat. Treatment right after shearing is most desirable. Repeat only as necessary. Do not use on dairy goats. See left label panel CAUTION for accepted dosage on young livestock; also other necessary livestock cautions. Do not treat animals under 3 months of age.

U.S.D.A. REG. NO. 606-79

MADE IN U.S.A.

ACCEPTED WITH COMMENTS

ACCEPTED

Jan. 8, 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC PEST CONTROL
ED UNDER NO. 606-79

[Handwritten signature]