

INSECT CONTROL: Flies: For control of flies in and around buildings which house domestic animals and around yards. 1 quart in 12 gallons of water. Apply or spray at rate of 1 gallon per 1,000 sq. ft. on painted surfaces and 2 gallons per 1,000 sq. ft. on unpainted surfaces where flies alight or congregate, such as walls, ceilings, stanchions, windows, fences around garbage cans, etc. Repeat applications as necessary. **CAUTION: Remove Lactating Dairy Animals from Buildings Before Spraying. Remove Animals Under One Month of Age Before Treating. Do Not Use Concentrations for Fly Control in Buildings on Animals.**

Mosquitoes: For an area or patio spray, add 1 quart to 7 gallons of water. Repeat applications as necessary.

CAUTION: AVOID PROLONGED BREATHING OF SPRAY MIST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN. WASH THOROUGHLY AFTER HANDLING. DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. MALATHION IS A CHOLINESTERASE INHIBITOR. DO NOT USE THIS PRODUCT ON ANIMALS SIMULTANEOUSLY OR WITHIN A FEW DAYS BEFORE OR AFTER TREATMENT WITH OR EXPOSURE TO CHOLINESTERASE INHIBITING DRUGS, PESTICIDES, OR CHEMICALS. ATROPINE IS ANTIDOTE. CONSULT VETERINARIAN AT FIRST SIGN OF ADVERSE REACTION. KEEP CONTAINER CLOSED. DO NOT USE POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

HIGHLY TOXIC TO FISH. DO NOT CONTAMINATE ANY BODY OF WATER BY DIRECT APPLICATION. CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES OR CONTAINERS.

USDA Reg. No. 606-96



MANGE & INSECT CONCENTRATE

A 57% Malathion concentrate to be diluted with water for application on farm animals and for fly and mosquito control.

ACTIVE INGREDIENTS

Malathion 57%

INERT INGREDIENTS

13% O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate

Caution: Keep out of reach of children. See other warnings on side panel.

NET CONTENTS

1 PINT

THE CORN KING Division
KING CASTLE, Inc.
 Marion, Iowa 52302

WARNING: APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL. Do Not Treat Animals Under One Month of Age. Do Not Contaminate Feed, Food Containers, or Watering Troughs. Do Not Apply to Lactating Dairy Animals or Non-Lactating Dairy Animals Within Two Weeks of Freshing.

DIRECTIONS

Use as a spray or dip diluted with water as follows:

BEEF CATTLE & HORSES: Lice: 1 quart in 25 gallons of water or 1 pint in 12½ gallons of water. Treat animals thoroughly. Repeat only if needed. **Ticks:** 1 to 2 quarts in 25 gallons of water or 1 to 2 pints in 12½ gallons of water. Treat animals thoroughly. Repeat application at two week intervals if needed.

Horn Fly: 1 to 1½ quarts in 25 gallons of water or 1 to 1½ pints in 12½ gallons of water. Apply complete coverage spray. Repeat at 2 week intervals if needed.

SWINE: Lice: 1 quart in 25 gallons of water or 1 pint in 12½ gallons of water. Treat animals, pens and litter thoroughly. Repeat only if needed.

Sarcoptic Mange: 1 quart in 25 gallons or 1 pint in 12½ gallons of water. Use extra care to clean all body surfaces including the inside of the ears. Repeat treatment in 10 days if needed. Further treatments may be made as needed but not more than once every two weeks. Animals should be kept out of sun and wind for 24 hours after treatment.

SHEEP AND GOATS: Lice, Ticks & Keds: 1 quart in 25 gallons of water or 1 pint in 12½ gallons of water. Treat animals thoroughly at application after 2 or 3 weeks if needed. **CAUTION: Do Not Apply to Milk Goats.**

See
 N
 10

