

Corn King

CATTLE SPRAY

DIRECTIONS (Continued)

for each 20 linear feet of bur-lap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

LICE CONTROL

(Dairy & Beef Cattle)

Make 1 to 2 applications of 3 to 3½ ounces of Corn King Cattle Spray at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

CAUTION

Do not treat animals within one day of slaughter. Do not apply dermally in combination with Trichlorofen. Harmful if swallowed. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not reuse container. Destroy when empty.

MADE IN U.S.A.

V1071

Corn King

CATTLE SPRAY

Protects Dairy and Beef Cattle against Horn, Face, Stable and House Flies.

Active Ingredients:

Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate*	1.00%
2, 2-dichlorovinyl dimethyl phosphate†	0.23%
Related compound†	0.02%
Petroleum Hydrocarbons	98.50%
Inert Ingredients	0.25%

* Ciodrin Insecticide
† Equivalent to 0.25% Vapona Insecticide

Caution: Keep out of reach of children. See other warnings on side panel.

**Net Contents
1 Gallon**

**Ciodrin
Vapona**
INSECTICIDES

Manufactured by
THE CORN KING Division
KING CASTLE, Inc.
Marion, Iowa 52302

Corn King

CATTLE SPRAY

DIRECTIONS

Corn King Cattle Spray is a specially prepared, ready-to-use insecticide suitable for use in conventional hand, power, or automatic spray equipment and in back rubbers. Apply with hand, power, or automatic equipment to moisten the hair only and not in excess of 2 fluid ounces daily per animal. Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under 6 months of age. Brahma cattle should not be treated as they may show hypersensitivity to organic phosphates. For the control of certain insects attacking animals, apply in accordance with State and Federal Recommendations.

FLY CONTROL

For the control of Horn Flies, Stable Flies, Face Flies, and House Flies on dairy and beef cattle apply as a light mist application at the rate of no more than 2 ounces daily. A 1 ounce application may be made twice daily or on dairy cows before morning and evening milking. Treat entire animal.

For Face Flies treat face and forehead.

For Stable Flies make sure spray reaches legs, ankles, and flanks.

BACK AND FACE RUBBERS

For the control of horn flies and face flies on dairy and beef animals use Corn King Cattle Spray in back rubbers and face rubbers. Ready to use; do not dilute. Use 1 gallon

E.P.A. Reg. No. 606-98