



DIRECTIONS (Continued)

Valuable in cage layer operations by sprinkling in the alleys and under cages. **Keep out of reach of birds.**

The number of applications per week is entirely dependent upon the fly population. It is important to always have the bait present in areas frequented by flies for constant control. Avoid sprinkling in litter and other places where it may sift away out of reach of flies.

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Induce vomiting if swallowed. Avoid contact with skin; wash thoroughly after using. Do not use inside homes. Do not contaminate milk, milk-handling equipment, feed or drinking water. If poisoning should occur, call a physician at once. Atropine and 2-PAM are the human antidotes.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.



FLY-BATE

CONTAINS DDVP and RONNEL

A Fly Bait for Use Either as it Comes from the Can or Mixed with Water and Used as a Syrup. Effective Against Flies Resistant to DDT or other Chlorinated Hydrocarbons.

Active Ingredients:

DDVP: 2, 2-dichlorovinyl dimethyl phosphate 0.32%
Related compounds 0.03%

Ronnel: 0-0-dimethyl 0-(2, 4, 5-trichlorophenyl)
phosphorothioate 0.25%

Inert Ingredients: 99.40%

Caution: Keep out of reach of children.
See other warnings on side panel.

Contents 1 Pound

Manufactured by

THE CORN KING Division
KING CASTLE, Inc.
Marion, Iowa 52302

DIRECTIONS

Use as a dry sprinkle fly bait as it comes from can, or dissolve 1 lb. in 1/2 gallon of water and use as a syrup. The syrup may be used in a sprinkler can or pieces of burlap may be saturated and hung up in areas frequented by flies. Make up only that amount of syrup needed for immediate use. Use wherever flies congregate, in all live-stock buildings including dairy barns and milk rooms, poultry houses, around garbage cans, dumps, and outside areas of human dwellings, restaurants, drive-ins, ice cream stands, dog kennels, etc. Sprinkle lightly either wet or dry on exposed surfaces such as floors, walkways, alleyways, etc. Use about 1/4 lb. per 1000 square feet of floor area. **Do not use in commercial food preparation areas or in edible product areas of food processing plants.**

E.P.A. Reg. No. 606-100

V372