



DIRECTIONS (Continued)

other openings. Do not use this solution for fogging or misting purposes in areas where animals have received a direct application within 8 hours. Do not contaminate water, feed or foodstuffs, milk or milking utensils.

**WARNING
Precautions In Using**

MAY BE FATAL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN. DO NOT BREATHE SPRAY MIST. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating and smoking. If swallowed induce vomiting immediately and get medical attention. Atropine is antidotal.

Decontaminate and destroy used containers. Do not re-use for any purpose. Decontaminate containers by rinsing thoroughly with water and detergent. Dispose of rinsings in a way as not to constitute a hazard or contaminate potable water supplies.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

© 1974 King Castle, Inc. All rights reserved.



**1% VAPONA[®]
ANIMAL & BARN
SPRAY**

For the Control of Horn Flies, Stable Flies, Face Flies, Mosquitoes and House Flies by Direct Application on Cows and Beef Cattle. For Control of Flies on and about Manure in Barns.

Active ingredients: 1% Vapona (Dieldrin) 1.00%
Inert ingredients 99.00%
Total 100.00%

Warning: Keep out of reach of children. See other warnings on side panel.

Net Contents

5 Gallons

Manufactured by
THE CORN KING Division
KING CASTLE, Inc.
Marion, Iowa 52302



DIRECTIONS FOR USE

1% Vapona[®] Spray is a specially prepared ready-to-use insecticide for use in conventional, hand, power, or automatic spray equipment, electric fogging or misting equipment.

Do not apply regularly to calves under 6 months of age. Erythromycin sulfate should not be treated if they may show hypersensitivity to the compound or phosphates.

Do not treat animals within 7 days of slaughter. Do not apply directly to the animal with Erythromycin.

HORN FLIES, STABLE FLIES, FACE FLIES, MOSQUITOES AND HOUSE FLIES—Apply 1% Vapona[®] Spray to the animal's body, especially the head and neck areas, by direct application. For control of flies on and about manure in barns, apply 1% Vapona[®] Spray to the manure. For control of flies on and about manure in barns, apply 1% Vapona[®] Spray to the manure. For control of flies on and about manure in barns, apply 1% Vapona[®] Spray to the manure.

For control of flies on and about manure in barns, apply 1% Vapona[®] Spray to the manure. For control of flies on and about manure in barns, apply 1% Vapona[®] Spray to the manure. For control of flies on and about manure in barns, apply 1% Vapona[®] Spray to the manure.