

301 / 5455222
18 / 201

16 DEC 1993

FILE COPY

Gail M. Dixon
Prince Agri Products, Inc.
P.O. Box 1009
Quincy, IL 62306

Dear Ms. Dixon:

Subject: Revised Labeling For:
Prince's Corn King Fly Spray
EPA Registration No. 606-103
Letter dated September 29, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

- 1) The statement, "It is a violation of Federal law to use this product in a manner inconsistent with its labeling." must appear on the label directly after the heading, "Directions For Use".
- 2) The following Statement Of Practical Treatment must appear on the label:

Statement Of Practical Treatment

If Swallowed: Call a physician or Poison Control Center immediately. Do not induce vomiting. Vomiting may cause aspiration pneumonia.

If Inhaled: Remove victim to fresh air.

If On Skin: Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes: Flush eyes with plenty of water. Call a physician immediately if irritation persists.

3) Add the following precautions to the label:


"Avoid skin and eye contact. Avoid breathing spray mist."

4) Add the following Environmental cautions under the heading "Environmental Hazards":

"This product is toxic to fish, birds, and other wildlife. Do not contaminate water when disposing of equipment washwater."

You may disregard our letter dated November 9, 1993. It was sent in error.

Sincerely,



Adam Heyward
Acting Product Manager 10
Insecticide & Rodenticide Branch
Registration Division (7505C)

CAUTION

KEEP OUT OF REACH OF CHILDREN. Do not spray near or toward an open flame. Avoid contamination of milk, milk utensils and feed. Harmful if swallowed.

Prevention Important— Effective control of flies requires the frequent removal of all manure, litter and other breeding places of flies.

Not for Household Use— To make the effect of Corn King's Fly Spray more lasting, an oil base is used. It is not recommended for household use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store near heat or open flame. Store container in upright position. Store in original container. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

E.P.A. Reg. No. 606-103
E.P.A. Est. No. 606-1A-1

ACCEPTED
with COMMENTS
in EPA Letter Dated

16 DEC 1993

TAKE TIME



Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

606-103

OBSERVE LABEL
DIRECTIONS

Corn King

FLY SPRAY

An oil base livestock spray with rapid knockdown and high killing powers.

For use on dairy cows, in dairy barns and on beef cattle and horses for control of face flies, horn flies, stable flies, house flies, horse flies, mosquitoes and gnats.

CAUTION:

Keep out of reach of children.
Observe additional cautions on back panel

Active Ingredients: Pyrethrins, 0.10%; piperonyl butoxide technical, 1.0%

Inert Ingredients: 98.9%**

* Equivalent to 0.80% of (butylcarbityl) (6-propylpiperonyl) ether and 0.20% of related compounds.

** Contains refined petroleum distillates

NET CONTENTS

2 x 2.50 Gallons
2 x 9.46 Liters

Manufactured by
PRINCE AGRI PRODUCTS, INC.
One Prince Plaza
Quincy, IL 62301

Corn King's Fly Spray applied to livestock with spray gun or insecticide applicator with considerable force is efficient. Make sure

On Dairy Cows: Do not spray during feeding. Do not spray efficiency spray closed during, and ing and spraying of

Using a Spray Gun: spray gun 15 to 20 each spraying use pint for each 12 to younger animals. cows and calves, animals thus helping and under parts soaked. Allow the face flies, spray head

Using an Insecticide: erly and operating tions as to the area; ger high enough so their feed is not c cows are exposed direct spraying. D may produce bronc Close windows a minutes fogging is

Spray Building: complete control of mosquitoes and gnats the animals and al may congregate. A liberally, directing minutes after spray

Spraying Beef Cattle: be used in the same buildings for killing