


[607-38]



U. S. D. A. Reg. No. 607 -- 38



Active Ingredients:	
Rotenone.....	0.75%
Other Cube Resins.....	2.00%
Inert Ingredients.....	97.25%
Net Wgt. 1 Lb.	100.00%

CCC GARDEN ROTE WITH ROTENONE

A PROVEN INSECTICIDE FOR VEGETABLES, FLOWERS AND FRUITS

Rotenone application is recognized generally as an effective control for many insect infestations. CCC contains no Paris Green, Lead or Calcium Arsenates. Contains no coarse particles. Economical. Sticks to foliage. Effective.

Apply with any dust gun under and top of leaves when leaves are moist if possible. Cover all infested portions of plants. Repeat when necessary to kill newly hatched insects. A little will go a long way. Be sure and use as directed. As a spray mix 1 lb. per 4 gals. water. Keep agitated. Do not apply to food crops within one day of harvest.

DIRECTIONS

BEANS, ASPARAGUS (Mexican Bean Beetle, Asparagus Beetle), **MELON, PUMPKIN, CUCUMBER, SQUASH** (Striped Beetle, Spotted Beetle, Aphis). Dust 15-20 lbs. per acre. Begin when eggs or beetles are found. Apply underside and tops of leaves. Cover plant thoroughly. Repeat as needed.

CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, BROCCOLI, KALE, KOHL RABI, TURNIPS (Cabbage Worm, Cabbage Looper, Flea Beetle, Caterpillars), **RADISH, BEETS, CARROTS, PARSNIPS, SWISS CHARD** (Webworms, Aphis), **STRAWBERRIES** (Leaf Roller, Spittle Bug), **TOMATO, EGG PLANT, PEPPER** (Young Tomato Hornworms, Colorado Potato Bug). Dust 20-25 lbs. per acre when insects appear. Repeat at 10-day intervals.

TOBACCO (Flea Beetles). Dust small plants 8-10 lbs. per acre. Large plants 15-25 lbs. per acre. Repeat every 7 days if necessary.

SPINACH, LETTUCE (Loopers, Aphis). Cover plant lightly. Dust 15-20 lbs. per acre.

POTATOES (Potato Beetles, Aphis). Cover plant thoroughly. Repeat as needed.

PEAS (Aphis). Most effective 7 days after application. 30 lbs. per acre. Repeat as needed. Dust blossoms and leaves. **JAPANESE BEETLE**, spray 1 lb. per 4 gals. water. Keep solution agitated. Spray foliage frequently to retain film during flying season of Beetles.

RASPBERRIES, BLACKBERRIES, GOOSEBERRIES (Raspberry Fruit Worm, Aphis). Dust 15 lbs. per acre. Spray 1 lb. to 4 gals. water (keep agitated). Last application 10 days before picking.

GRAPES, APPLES (Aphis), **CURRANTS** (Aphis). Cover plant thoroughly as a dust or 1 lb. to 4 gals. water as a spray (keep agitated). Repeat as needed.

NASTURTIUMS, ROSES, ETC. (Aphis). Dust thoroughly but lightly or spray 1 lb. to 4 gals. water (keep agitated). Repeat as needed.

THIS CAN CONTAINS FULL NET WGT. AS INDICATED. SOME SETTLING MAY OCCUR DURING SHIPPING AND HANDLING.

Carbola Chemical Co., Inc., Natural Bridge, N.Y., U.S.A.

PRINTED IN U.S.A.

