

10/10/68 with 11/5/68 deletion per USDA 11/4/68. BK

Labeling label  
(CAREOLA SPECIAL DAIRY CATTLE DUST)  
Carbola Chemical Company, Inc.  
Natural Bridge, N. Y. 13665

CONTENTS PANEL

Copy A: CAREOLA SPECIAL DAIRY CATTLE DUST (1 lb.)

Copy B: For Dairy and Beef Cattle

Copy C: Contains Malathion and Methoxychlor

Copy D: Controls Hornflies on Dairy Cattle and Hornflies,  
Lice and Ticks on Beef Cattle

Copy E: Active Ingredients:

Methoxychlor, technical<sup>®</sup>..... 5.0  
Malathion (S, C-dimethyl dithiophosphate of  
methyl mercaptosuccinate)..... 4.0  
Inert Ingredients:..... 91.0

<sup>®</sup>Equivalent to 4.0% 2,2-Dim (p-methoxyphenyl)-1,  
1,1-trichloroethane and 0.6% related reaction  
products.

Copy F: USDA Reg. No. \_\_\_\_\_

Copy G: CAUTION: KEEP OUT OF REACH OF CHILDREN.  
See back panel for additional cautions.

Copy H: CARBOLA CHEMICAL COMPANY, INC.  
A subsidiary of International Vale Co., Inc.  
Natural Bridge, N. Y. 13665

ACCEPTED  
NOV 25 1968  
UNDER THE FEDERAL  
REGISTRATION AND  
FOR FRODOX  
ED UNDER NO. 601-43  
TO ATTACHED COMMENTS

10/10/68

with 11/5/68 deletion per USDA 11/4/68 KR

Labeling Label  
SPECIAL DAIRY CATTLE DUST  
Parke-Davis Chemical Co., Inc.  
New York, N. Y. 10013

Copy I: FOR DAIRY AND BEEF CATTLE

Copy J: DIRECTIONS FOR USE

FOR DIRECTIONS OF APPLICATION TO DAIRY CATTLE AND BEEF CATTLE, FOR CONTROL OF HORNETS:  
Apply 2 heaping tablespoonfuls (2 ounces) to the backs and flanks of infested animals. Rub lightly into hair. Repeat application no more than once in every 3-week period, if necessary.

FOR CONTROL OF LICE AND TICKS ON BEEF CATTLE:  
Use as above. Treat animals thoroughly. Repeat treatments as necessary.

AS AN AID FOR THE CONTROL OF FACE FLIES ON BEEF CATTLE: Dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.

FOR CONTROL OF HORSE LICE: Apply to legs, pasterns, and hocks. This treatment may be sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed.

LICE CONTROL ON SHEEP: Treat animals thoroughly, preferably following shearing. Repeat as necessary.

~~FOR CONTROL OF HORNETS ON SHEEP: Apply to the neck and head of infested animals. Repeat as necessary.~~

DELETED  
11/6/68  
K. ROAST

Copy K: CAUTION: Harmful if swallowed. Do not apply within 5 hours of milking or during milking. Do not contact with feed or water. Avoid prolonged breathing of dust. Avoid prolonged contact with skin; wash thoroughly after using. Avoid contamination of feed and feed water.