


Wormer

THIABENDAZOLE
HORSE

CONTROLS BOTS AND ROUNDWORMS



NET WT. 0.42 OZ.

TRICHLORFON
WORMER




both compartments

TO OPEN CUT

TOP FORM Wormer Two*

(THIABENDAZOLE
HORSE

CONTROLS BOTS AND ROUNDWORMS



OR TEAR HERE

TRICHLORFON)
WORMER

EPA REG. NO. 618-85-AA

TO OPEN CUT

TOP FORM WORMER TWO

For Oral Use in Horses.

Provides control of bots and roundworms in one treatment.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

USE DIRECTIONS: Use one package (0.42 OZ.) for each 250 lb. body weight.

Tear off the top of the package so that the entire contents of both compartments can be emptied. Sprinkle the entire contents of the package on a portion of the usual daily ration. Treat each horse individually.

Most horses find TOP FORM WORMER TWO acceptable. Allow fussy eaters more time to consume the medicated feed. Allow horses to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed. For the horse that refuses to eat medicated feed, consult your veterinarian.

RECOMMENDED WORM CONTROL PROGRAM: Many horses need regular worming for bloodworms and other roundworms every 6 to 8 weeks. TOP FORM WORMER TWO may be used for worming throughout the year. For the most effective control of bots, treatment is recommended approximately one month following a killing frost.

*Trademark

†U.S. Pat. 3,017,415

**Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poterostomum*, *Oxyuris*, *Strongyloides*, *Parascaris*, and *Gasterophilus*.

OR TEAR

E

CONTRA
debilitate
age, mar
or anima
minister
cially mu
weeks al
be slaug
Warning
animals
days be
exposur
drugs, p
NOTICE
over-dos
ataxia at
100 mg.
weight, i
PRECAU
mended
more fre
WARNIN
contact
soap and
humans,
tablespo
Call a p
CIAN: At
Give sup
patient

TC
MERCK
MERCK & C
Store bel
Lot No. 8
Exp. Date

ACCEPTED

MAR 27 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 618-85

8 23 74

1/22/75

OR TEAR HERE

TRICHLORFON WORMER

EPA REG. NO. 618-85-AA

TO OPEN CUT

TOP FORM WORMER TWO

For Oral Use in Horses.

Provides control of bots and roundworms in one treatment.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

USE DIRECTIONS: Use one package (0.42 OZ) for each 250 lb. body weight.

Tear off the top of the package so that the entire contents of both compartments can be emptied. Sprinkle the entire contents of the package on a portion of the usual daily ration. Treat each horse individually.

Most horses find TOP FORM WORMER TWO acceptable. Allow fussy eaters more time to consume the medicated feed. Allow horses to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed. For the horse that refuses to eat medicated feed, consult your veterinarian.

RECOMMENDED WORM CONTROL PROGRAM: Many horses need regular worming for bloodworms and other roundworms every 6 to 8 weeks. TOP FORM WORMER TWO may be used for worming throughout the year. For the most effective control of bots, treatment is recommended approximately one month following a killing frost.

*Trademark

†U.S. Pat. 3,017,415

**Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, *Strongyloides*, *Parascaris*, and *Gasterophilus*.

OR TEAR HERE

EPA Est. 10036-NJ-1

Use Only As Directed

CONTRAINDICATIONS: Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using. Do not use in horses to be slaughtered for food purposes.

Warning—Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

NOTICE TO VETERINARIAN: Symptoms of over-dosage of trichlorfon in horses are ataxia and colic. Atropine, at the rate of 100 mg. (1 1/2 grains) per 100 lb. body weight, is an antidote.

PRECAUTIONS: Do not exceed recommended dose. Do not repeat treatment more frequently than every 30 days.

WARNING: Avoid prolonged or repeated contact with the skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in a glass of water. Call a physician immediately. TO PHYSICIAN: Atropine and 2-PAM are antidotes. Give supportive treatment. Do not walk patient or give morphine.

Product 52341

TOP FORM PRODUCTS

MERCK CHEMICAL DIVISION

MERCK & CO., INC. • RAHWAY, N.J. 07065, U.S.A.

Store below 86°F (30°C)

Lot No. &
Exp. Date

ACCEPTED

MAR 27 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 618-85

8 23 74

1/22/75

8 23 74

1/22/75

TO OPEN CUT

TOP FORM WORMER TWO

For Oral Use in Horses.

Provides control of bots and roundworms in one treatment.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

USE DIRECTIONS: Use one package (0.42 OZ.) for each 250 lb. body weight.

Tear off the top of the package so that the entire contents of **both compartments** can be emptied. Sprinkle the entire contents of the package on a portion of the usual daily ration. Treat each horse individually.

Most horses find TOP FORM WORMER TWO acceptable. Allow fussy eaters more time to consume the medicated feed. Allow horses to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed. For the horse that refuses to eat medicated feed, consult your veterinarian.

RECOMMENDED WORM CONTROL PROGRAM: Many horses need regular worming for bloodworms and other roundworms every 6 to 8 weeks. TOP FORM WORMER TWO may be used for worming throughout the year. For the most effective control of bots, treatment is recommended approximately one month following a killing frost.

*Trademark

†U.S. Pat. 3,017,415

**Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, *Strongyloides*, *Parascaris*, and *Gasterophilus*.

OR TEAR HERE

EPA Est. 10036-NJ-1

Use Only As Directed

CONTRAINDICATIONS: Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using. Do not use in horses to be slaughtered for food purposes.

Warning—Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

NOTICE TO VETERINARIAN: Symptoms of over-dosage of trichlorfon in horses are ataxia and colic. Atropine, at the rate of 100 mg. (1 1/2 grains) per 100 lb. body weight, is an antidote.

PRECAUTIONS: Do not exceed recommended dose. Do not repeat treatment more frequently than every 30 days.

WARNING: Avoid prolonged or repeated contact with the skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in a glass of water. Call a physician immediately. TO PHYSICIAN: Atropine and 2-PAM are antidotes. Give supportive treatment. Do not walk patient or give morphine.

Product 52341

TOP FORM PRODUCTS

MERCK CHEMICAL DIVISION

MERCK & CO., INC. • RAHWAY, N.J. 07065, U.S.A.

Store below 86°F (30°C)

Lot No. &

Exp. Date


ACCEPTED

MAR 27 1975

TO OPEN:

**TOP FORM
Wormer
Two***
(THIABENDAZOLE
HORSE

CONTROLS BOTS AND ROUNDWORMS




CUT OR TEAR HERE

TRICHLORFON)
WORMER

Wormer

(THIABENDAZOLE
HORSE

CONTROLS BOTS AND ROUNDWORMS



TRICHLORFON
WORMER

For the control of bots, large strongyles, small strongyles, pinworms, ascarids, and ascarids in horses.** See back panel for use directions.

This package will treat one 1000 lb. horse.

For larger or smaller horses, a 0.42 oz. package is available (treats 250 lb. body weight).

This package contains
20 g. Thiabendazole† and 18 g. Trichlorfon

WARNING!

Keep out of reach of children.

See back panel for other precautions.

EPA REG. NO. 618-85-AA EPA Est. 10036-NJ-1

TO OPEN CUT

TOP FORM WORMER TWO

For Oral Use in Horses.

Provides control of bots and roundworms in one treatment.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

USE DIRECTIONS: Use one package (1 $\frac{2}{3}$ oz.) for each 1000 lb. horse. For larger or smaller horses, 0.42 oz. package is available (treats 250 lb. body weight).

Tear off the top of the package so that the entire contents of **both compartments** can be emptied. Sprinkle the entire contents of the package on a portion of the usual daily ration. Treat each horse individually.

Most horses find TOP FORM WORMER TWO acceptable. Allow fussy eaters more time to consume the medicated feed. Allow horses to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed. For the horse that refuses to eat medicated feed, consult your veterinarian.

RECOMMENDED WORM CONTROL PROGRAM: Many horses need regular worming for bloodworms and other roundworms every 6 to 8 weeks. TOP FORM WORMER TWO may be used for worming throughout the year. For the most effective control of bots, treatment is recommended approximately one month following a killing frost.

*Trademark

†U.S. Pat. 3,017,415

**Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, *Strongylodontes*, *Parascaris*, and *Gasterophilus*.

OR TEAR HERE

Use Only As Directed

CONTRAINDICATIONS: Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using. Do not use in horses to be slaughtered for food purposes.

Warning—Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

NOTICE TO VETERINARIAN: Symptoms of overdosage of trichlorfon in horses are ataxia and colic. Atropine, at the rate of 100 mg. (1 $\frac{1}{2}$ grains) per 100 lb. body weight, is an antidote.

PRECAUTIONS: Do not exceed recommended dose. Do not repeat treatment more frequently than every 30 days.

WARNING: Avoid prolonged or repeated contact with the skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in a glass of water. Call a physician immediately. TO PHYSICIAN: Atropine and 2-PAM are antidotes. Give supportive treatment. Do not walk patient or give morphine.

Product **52341**

TOP FORM PRODUCTS

MERCK CHEMICAL DIVISION

MERCK & CO. INC. • RAHWAY, N.J. 07065, U.S.A.

Store below 86°F (30°C)

Lot No. &
Exp. Date



TOP FORM WORMER TWO

For Oral Use in Horses.

Provides control of bots and roundworms in one treatment.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

USE DIRECTIONS: Use one package (1½ oz.) for each 1000 lb. horse. For larger or smaller horses, 0.42 oz. package is available (treats 250 lb. body weight).

Tear off the top of the package so that the entire contents of **both compartments** can be emptied. Sprinkle the entire contents of the package on a portion of the usual daily ration. Treat each horse individually.

Most horses find TOP FORM WORMER TWO acceptable. Allow fussy eaters more time to consume the medicated feed. Allow horses to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed. For the horse that refuses to eat medicated feed, consult your veterinarian.

RECOMMENDED WORM CONTROL PROGRAM: Many horses need regular worming for bloodworms and other roundworms every 6 to 8 weeks. TOP FORM WORMER TWO may be used for worming throughout the year. For the most effective control of bots, treatment is recommended approximately one month following a killing frost.

Use Only As Directed

CONTRAINDICATIONS: Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using. Do not use in horses to be slaughtered for food purposes.

Warning — Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

NOTICE TO VETERINARIAN: Symptoms of over-dosage of trichlorfon in horses are ataxia and colic. Atropine, at the rate of 100 mg. (1½ grains) per 100 lb. body weight, is an antidote.

PRECAUTIONS: Do not exceed recommended dose. Do not repeat treatment more frequently than every 30 days.

WARNING: Avoid prolonged or repeated contact with the skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in a glass of water. Call a physician immediately. TO PHYSICIAN: Atropine and 2-PAM are antidotes. Give supportive treatment. Do not walk patient or give morphine.

*Trademark

†U.S. Pat. 3,017,415

**Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craierostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, *Strongyloides*, *Parascaris*, and *Gasterophilus*.

Product 52341

TOP FORM PRODUCTS

MERCK CHEMICAL DIVISION

MERCK & CO., INC. • RAHWAY, N.J. 07065, U.S.A.

Store below 86°F (30°C)

Lot No. &
Exp. Date ►

TOP FORM Wormer

TWO*

(THIABENDAZOLE—TRICHLORFON)

HORSE WORMER

CONTROLS BOTS AND ROUNDWORMS



TOP FIRM WORMER TWO

Departments


both.com

Use Only As Directed

52411
M
MERCK CHEMICAL DIVISION
KILGORE, TEXAS

Wormer

(THIABENDAZOLE—TRICHLORFON)
HORSE WORMER
CONTROLS BOTS AND ROUNDWORMS



EPA REG. NO. 618-85-AA

EPA Est. 10036-NJ-1

For the control of bots, large strongyles, small strongyles, pinworms, *Strongyloides*, and ascarids in horses.** See back panel for use directions.

This package will treat one 1000 lb. horse.

For larger or smaller horses, a 0.42 oz. package is available (treats 250 lb. body weight).

This package contains 20 g. Thiabendazole† and 18 g. Trichlorfon.

WARNING!

Keep out of reach of children. See back panel for other precautions.

NET WT. 1 2/3 OZ.