

TO OPEN CUT

OR TEAR HERE

Product 65701

**Equizole**


**(THIABENDAZOLE**

**HORSE**

**B**

**TRICHLORFON)**

**WORMER**



**CAUTION** Federal law restricts this drug to use by or on the order of a licensed veterinarian

This package contains 5 g Thiabendazole and 4.5 g Trichlorfon

See back panel for use directions and precautions

Store below 86° F (30° C)

† U.S. Pat. 3,017,415

EPA REG. NO. 618-88-AA

EPA Est. 10036-NJ-1

Professional Veterinary Products  
**MERCK CHEMICAL DIVISION**  
MERCK & CO., INC. • RAHWAY, N.J. 07065 U.S.A.

TO OPEN CUT

OR T

**INDICATIONS:** For the control of large strongyles, small strongyles, pinworms, *Strongyloides*, ascarids, and bots in horses.\*\*

**DOSAGE:** This package (0.42 oz.) will treat 250 lb. of body weight.

**USE DIRECTIONS:** For a satisfactory diagnosis, a microscopic fecal examination should be performed prior to worming.

**EQUIZOLE B**, at the recommended dose rate of 2 g. of thiabendazole and 1.8 g. of trichlorfon per 100 lb. of body weight, is highly effective against all genera listed below\*\* It may be given by stomach tube, drench, or in feed

**Stomach Tube or Drench**— Tear off top of package so that entire contents of **both compartments** are emptied into a suitable container. Add 5 to 10 fl. oz. of water. Stir until thoroughly mixed and give by stomach tube or drench. This suspension may be stored for 24 hours. Prepare doses individually.

**Feed**— Sprinkle the entire contents of **both compartments** of the package on a portion of the usual daily ration. Allow horse to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed.

**NOTE:** Some horses are quite selective about their rations and do not readily accept changes or new ingredients in their diet. A few hours fast or a moist feed containing molasses or sweetened with sugar may help improve acceptability in such animals.

\*\*Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, *Strongyloides*, *Parascaris*, and *Gasterophilus*

CO  
del  
of  
na  
no  
esj  
of  
in  
po  
on  
da  
po  
pe  
NC  
ov  
at:  
10  
we  
PR  
me  
rel  
ev  
Wi  
co  
so  
hu  
sp  
a  
At  
su  
or

L  
E

5/3/74

1/22/75

OPEN CUT

OR TEAR HERE

Product 65701

B

TRICHLORFON)

WORMER



EPA REG. NO. 618-88-AA

EPA Est. 10036-NJ-1

Professional Veterinary Products  
MERCK CHEMICAL DIVISION  
MERCK & CO., INC. • RAHWAY, N. J. 07065, U.S.A.

TO OPEN CUT

OR TEAR HERE

**INDICATIONS:** For the control of large strongyles, small strongyles, pinworms, *Strongyloides*, ascarids, and bots in horses.\*\*

**DOSAGE:** This package (0.42 oz.) will treat 250 lb. of body weight.

**USE DIRECTIONS:** For a satisfactory diagnosis, a microscopic fecal examination should be performed prior to worming.

EQUIZOLE B, at the recommended dose rate of 2 g. of thiabendazole and 1.8 g. of trichlorfon per 100 lb. of body weight, is highly effective against all genera listed below.\*\* It may be given by stomach tube, drench, or in feed.

**Stomach Tube or Drench**—Tear off top of package so that entire contents of **both compartments** are emptied into a suitable container. Add 5 to 10 fl. oz. of water. Stir until thoroughly mixed and give by stomach tube or drench. This suspension may be stored for 24 hours. Prepare doses individually.

**Feed**—Sprinkle the entire contents of **both compartments** of the package on a portion of the usual daily ration. Allow horse to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours after being offered. Not all horses will consume medicated feed.

**NOTE:** Some horses are quite selective about their rations and do not readily accept changes or new ingredients in their diet. A few hours fast or a moist feed containing molasses or sweetened with sugar may help improve acceptability in such animals.

\*\*Including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, *Strongyloides*, *Parascaris*, and *Gasterophilus*.

**Use Only As Directed**

**CONTRAINDICATIONS:** Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using. Do not use in horses to be slaughtered for food purposes. **Warning**—Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

**NOTICE TO VETERINARIAN:** Symptoms of over-dosage of trichlorfon in horses are ataxia and colic. Atropine, at the rate of 100 mg. (1½ grains) per 100 lb. body weight, is an antidote.

**PRECAUTIONS:** Do not exceed recommended dose. Not for human use. Do not repeat treatment more frequently than every 30 days.

**WARNING:** Avoid prolonged or repeated contact with the skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in a glass of water. Call a physician immediately. **TO PHYSICIAN:** Atropine and 2 PAM are antidotes. Give supportive treatment. Do not walk patient or give morphine.

KEEP OUT OF REACH OF CHILDREN

Lot No & Exp. Date ▶

5/3/74

1/22/75

5/3/74

TO OPEN. CUT

OR TEAR HERE

Product 65701

**THIABENDAZOLE**

**B**

**(THIABENDAZOLE — TRICHLORFON)**

**HORSE**

**WORMER**



CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

This package contains 20 g. Thiabendazole and 18 g. Trichlorfon

See back panel for use directions and precautions

Store below 86° F (30° C)

† U.S. Pat. 3,017,415

EPA REG. NO. 618-86-AA

EPA Est. 10036-NJ-1

Professional Veterinary Products  
**MERCK CHEMICAL DIVISION**  
MERCK & CO., INC. - RAHWAY, N.J. 07065 U.S.A.